

FILE 'MEDLINE' ENTERED AT 18:34:51 ON 29 DEC 2002

FILE 'CAPLUS' ENTERED AT 18:34:51 ON 29 DEC 2002  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'LIFESCI' ENTERED AT 18:34:51 ON 29 DEC 2002  
COPYRIGHT (C) 2002 Cambridge Scientific Abstracts (CSA)

FILE 'EMBASE' ENTERED AT 18:34:51 ON 29 DEC 2002  
COPYRIGHT (C) 2002 Elsevier Science B.V. All rights reserved.

FILE 'USPATFULL' ENTERED AT 18:34:51 ON 29 DEC 2002  
CA INDEXING COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'BIOSIS' ENTERED AT 18:34:51 ON 29 DEC 2002  
COPYRIGHT (C) 2002 BIOLOGICAL ABSTRACTS INC.(R)

=> s hiv and tar  
L4 3924 HIV AND TAR

=> s l4 and snorna  
L5 1 L4 AND SNORNA

=> d l5 ibib abs tot

L5 ANSWER 1 OF 1 USPATFULL

ACCESSION NUMBER: 1999:81758 USPATFULL  
TITLE: Non-activated receptor complex proteins and uses thereof  
INVENTOR(S): Davis, Roger J., Princeton, MA, United States  
Galcheva-Gargova, Zoya, Worcester, MA, United States  
PATENT ASSIGNEE(S): University of Massachusetts, Boston, MA, United States (U.S. corporation)

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 5925566		19990720
APPLICATION INFO.:	US 1997-870518		19970606 (8)

	NUMBER	DATE
PRIORITY INFORMATION:	US 1996-19219P	19960606 (60)
DOCUMENT TYPE:	Utility	
FILE SEGMENT:	Granted	
PRIMARY EXAMINER:	Campbell, Bruce R.	
ASSISTANT EXAMINER:	Nguyen, Dave Trong	
LEGAL REPRESENTATIVE:	Fish & Richardson, P.C.	
NUMBER OF CLAIMS:	23	
EXEMPLARY CLAIM:	1	
NUMBER OF DRAWINGS:	22 Drawing Figure(s); 18 Drawing Page(s)	
LINE COUNT:	2438	

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB The invention features a substantially pure ZPR1 polypeptide. For example, a ZPR1 polypeptide that specifically binds to a non-activated membrane-bound receptor (e.g., EGF or PDGF receptors) and specifically binds small nucleolar RNAs (e.g., U3). ZPR1 polypeptides can be isolated from any eukaryote, including mammals (e.g. rodents and humans) and fungi (e.g., *S. cerevisiae* and *S. pombe*).

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

=> d kwic

L5 ANSWER 1 OF 1 USPATFULL

SUMM "Small nucleolar RNA" or " \*\*\*snRNA\*\*\* " refers to RNA sequences located in the nucleolus and includes the sequences encoded by the U3, U10, U15, U22, and . . .  
DETD . . . The interaction of ZPR1 with the snRNAs U3, U10, U15, U22, U33, or the viral RNAs corresponding to the RRE ( \*\*\*HIV\*\*\* Rev responsive element), \*\*\*HIV\*\*\* trans-acting response element ( \*\*\*TAR\*\*\* ), and adenovirus VA-RNA1 (Akusjavari et al., Proc. Natl.

. (1988); Bartel. . . was investigated by this method. No binding of ZPR1 was detected in experiments using viral RNA corresponding to RRE, the \*\*\*TAR\*\*\*, or adenovirus VA-RNA1. In contrast, binding to ZPR1 was detected in experiments using several small nucleolar RNAs (snRNAs), including U3, . . .

DETD The specificity of the interaction of RNA with ZPR1 was examined in greater detail in competition experiments using the \*\*\*snRNA\*\*\* U3. A [.sup.32 P]-labeled U3 probe was incubated with ZPR1 alone or in the presence of excess non-radioactive U3, anti-sense. . .

=>  
=> d history

FILE 'MEDLINE, CAPLUS, LIFESCI, EMBASE, USPATFULL, BIOSIS' ENTERED AT  
18:34:51 ON 29 DEC 2002

L4 3924 S HIV AND TAR  
L5 1 S L4 AND SNORNA

=> s l4 and u16  
L6 1 L4 AND U16

=> s l6 not l5  
L7 1 L6 NOT L5

=> d l7 ibib abs tot

L7 ANSWER 1 OF 1 USPATFULL  
ACCESSION NUMBER: 2001:142079 USPATFULL  
TITLE: Method for screening nucleic acid catalysts  
INVENTOR(S): Burgin, Alex, Chula Vista, CA, United States  
Beigelman, Leonid, Longmont, CO, United States  
Bellon, Laurent, Boulder, CO, United States  
PATENT ASSIGNEE(S): Ribozyne Pharmaceuticals, Inc., Boulder, CO, United States (U.S. corporation)

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 6280936	B1	20010828
APPLICATION INFO.:	US 1998-94381		19980609 (9)

	NUMBER	DATE
PRIORITY INFORMATION:	US 1997-49002P	19970609 (60)
DOCUMENT TYPE:	Utility	
FILE SEGMENT:	GRANTED	
PRIMARY EXAMINER:	Fredman, Jeffrey	
LEGAL REPRESENTATIVE:	McDonnell Boehnen Hulbert & Berghoff	
NUMBER OF CLAIMS:	24	
EXEMPLARY CLAIM:	1	
NUMBER OF DRAWINGS:	23 Drawing Figure(s); 23 Drawing Page(s)	
LINE COUNT:	2403	

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB Nucleic acid catalysts, method of screening for variants of nucleic acid catalysts, synthesis of ribozyme libraries and discovery of gene sequences are described.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

=> d kwic

L7 ANSWER 1 OF 1 USPATFULL  
DETD In yet another embodiment, the nucleotide linker (L) is a nucleic acid aptamer, such as an ATP aptamer, \*\*\*HIV\*\*\* Rev aptamer (RRE), \*\*\*HIV\*\*\* Tat aptamer ( \*\*\*TAR\*\*\* ) and others (for a review see Gold et al., 1995, Annu. Rev. Biochem., 64, 763; and Szostak & Ellington, 1993, . . .

DETD . . . that for the hammerhead ribozyme one residue (A15.1) must remain constant; A15.1 forms a base pair with a substrate nucleotide ( \*\*\*U16\*\*\* .1) but is also absolutely required for ribozyme activity. It is the only residue within the hammerhead ribozyme that is part. . .

E1	.	2	MICHIELUTTI RONDA E/AU
E2		5	MICHIELUTTI THOMAS J/AU
E3		0 -->	MICHIEZENI ?/AU
E4		27	MICHIEZENI A/AU
E5		21	MICHIEZENI ALESSANDRO/AU
E6		2	MICHIEZENI F/AU
E7		3	MICHIEZENI GIACOMO F/AU
E8		6	MICHIEZENI K E/AU
E9		1	MICHIEZENI KATHLEEN E/AU
E10		3	MICHIEZENI L J/AU
E11		15	MICHIEZENI M/AU
E12		1	MICHIEZENI M A/AU

=> s e4-5  
L8 48 ("MICHIEZENI A"/AU OR "MICHIEZENI ALESSANDRO"/AU)

=> dup rem l8  
PROCESSING COMPLETED FOR L8  
L9 15 DUP REM L8 (33 DUPLICATES REMOVED)

=> d l9 ibib abs tot

L9 ANSWER 1 OF 15 MEDLINE DUPLICATE 1  
ACCESSION NUMBER: 2002647867 MEDLINE  
DOCUMENT NUMBER: 22294961 PubMed ID: 12376617  
TITLE: A nucleolar TAR decoy inhibitor of HIV-1 replication.  
AUTHOR: \*\*\*Michienzi Alessandro\*\*\* ; Li Shirley; Zaia John A;  
Rossi John J  
CORPORATE SOURCE: Divisions of Molecular Biology and Virology, Beckman  
Research Institute of the City of Hope, 1450 East Duarte  
Road, Duarte, CA 91010-3011, USA.  
CONTRACT NUMBER: AI29329 (NIAID)  
AI46030 (NIAID)  
SOURCE: PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE  
UNITED STATES OF AMERICA, (2002 Oct 29) 99 (22) 14047-52.  
Journal code: 7505876. ISSN: 0027-8424.  
PUB. COUNTRY: United States  
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 200212  
ENTRY DATE: Entered STN: 20021031  
Last Updated on STN: 20021217  
Entered Medline: 20021209

AB Tat is a critical regulatory factor in HIV-1 gene expression. It mediates  
the transactivation of transcription from the HIV-1 LTR by binding to the  
transactivation response (TAR) element in a complex with cyclin T1.  
Because of its critical and early role in HIV gene expression, Tat and its  
interaction with the TAR element constitute important therapeutic targets  
for the treatment of HIV-1 infection. Based on the known nucleolar  
localization properties of Tat, we constructed a chimeric small nucleolar  
RNA-TAR decoy that localizes to the nucleoli of human cells and  
colocalizes in the nucleolus with a Tat-enhanced GFP fusion protein. When  
the chimeric RNA was stably expressed in human T lymphoblastoid CEM cells  
it potently inhibited HIV-1 replication. These results demonstrate that  
the nucleolar trafficking of Tat is critical for HIV-1 replication and  
suggests a role for the nucleolus in HIV-1 viral replication.

L9 ANSWER 2 OF 15 CAPLUS COPYRIGHT 2002 ACS  
ACCESSION NUMBER: 2002:899156 CAPLUS  
TITLE: Intracellular ribozyme applications  
AUTHOR(S): Castanotto, D.; Li, J. R.; \*\*\*Michienzi, A.\*\*\* ;  
Langlois, M.-A.; Lee, N. S.; Puymirat, J.; Rossi, J.  
J.  
CORPORATE SOURCE: Division of Molecular Biology, Beckman Research  
Institute of the City of Hope, Duarte, CA, 91010-3011  
SOURCE: Biochemical Society Transactions (2002), 30(6),  
1140-1145  
CODEN: BCSTB5; ISSN: 0300-5127  
PUBLISHER: Portland Press Ltd.  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB The exquisite target selectivity of trans -acting ribozymes has fostered  
their use as potential therapeutic agents and tools for down-regulating  
cellular transcripts. In living cells, free diffusion of RNAs is  
extremely limited, if it exists at all. Thus, getting ribozymes to

- transcript with the target RNA. In addn., not all sites along a given target RNA are equally accessible to ribozyme base pairing. Cellular proteins greatly influence the trafficking and structure of RNA, and therefore making ribozymes work effectively in cells a significant challenge. This article addresses the problems of getting engineered ribozymes to effectively pair with and cleave targets in cells. The work described here illuminates methods for target-site selection on native mRNAs, methods for ribozyme expression, and strategies for obtaining a discrete intracellular localization of ribozymes.

L9 ANSWER 3 OF 15 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.  
 ACCESSION NUMBER: 2002:624770 BIOSIS  
 DOCUMENT NUMBER: PREV200200624770  
 TITLE: Anti-HIV therapy: New concepts & results retroviral delivery of combinations of anti-HIV RNAs in cord blood CD34+ and HOS-CCR5/CD4 cells.  
 AUTHOR(S): Kleihauer, Annette (1); Kim, James (1); Bauer, Gerhard (1); Cagnon, Laurance (1); \*\*\*Michienzi, Alessandro (1)\*\*\* ; Zaia, John A. (1); Rossi, John J. (1)  
 CORPORATE SOURCE: (1) City of Hope National Medical Center/Beckman Research Institute, Duarte, CA, 91010 USA  
 SOURCE: Journal of Human Virology, (January February, 2002) Vol. 5, No. 1, pp. 89-90. <http://www.humanvirology.com/>. print. Meeting Info.: 2002 International Meeting of the Institute of Human Virology Baltimore, Maryland, USA September 09-13, 2002 Institute of Human Virology . ISSN: 1090-9508.  
 DOCUMENT TYPE: Conference  
 LANGUAGE: English

L9 ANSWER 4 OF 15 CAPLUS COPYRIGHT 2002 ACS  
 ACCESSION NUMBER: 2002:872767 CAPLUS  
 TITLE: Chimeric snRNA-Rev binding aptamer RNA as Rev protein decoy and inhibitor of HIV1 replication  
 INVENTOR(S): Bozzoni, Irene; \*\*\*Michienzi, Alessandro\*\*\* ; Buonomo, Sara Cristina Barbara  
 PATENT ASSIGNEE(S): Universita' Degli Studi Di Roma 'la Sapienza', Italy  
 SOURCE: Ital., 35 pp. CODEN: ITXXBY  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Italian  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
IT 1305300	B1	20010504	IT 1999-RM126	19990224
PRIORITY APPLN. INFO.:			IT 1999-RM126	19990224

AB A chimeric RNA mol. which functions as a decoy for HIV1 Rev protein is disclosed. This chimeric RNA consists of an snRNA fragment, which is responsible for translocation of the chimera to the nucleolus, and a Rev protein-binding RNA fragment. Thus, a construct encoding an snRNA U16 fragment fused to the Rev binding element of RRE was prepd. In recombinant 293 cells the chimeric RNA was found in the nucleus where it bound to Rev protein and was exported to the cytoplasm.

L9 ANSWER 5 OF 15 MEDLINE  
 ACCESSION NUMBER: 2002680695 IN-PROCESS  
 DOCUMENT NUMBER: 22328739 PubMed ID: 12440991  
 TITLE: Intracellular ribozyme applications.  
 AUTHOR: Castanotto D; Li J R; \*\*\*Michienzi A\*\*\* ; Langlois M A; Lee N S; Puymirat J; Rossi J J  
 CORPORATE SOURCE: Division of Molecular Biology, Beckman Research Institute of the City of Hope, Duarte, CA 91010-3011, U.S.A.  
 SOURCE: BIOCHEMICAL SOCIETY TRANSACTIONS, (2001 Dec) 30 (6) 1140-5. Journal code: 7506897. ISSN: 0300-5127.  
 PUB. COUNTRY: England; United Kingdom  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: IN-PROCESS; NONINDEXED; Priority Journals  
 ENTRY DATE: Entered STN: 20021121  
 Last Updated on STN: 20021213

AB The exquisite target selectivity of trans -acting ribozymes has fostered their use as potential therapeutic agents and tools for down-regulating cellular transcripts. In living cells, free diffusion of RNAs is extremely limited, if it exists at all. Thus, getting ribozymes to base-pair with

- the target RNA. In addition, not all sites along a given target RNA are equally accessible to ribozyme base pairing. Cellular proteins greatly influence the trafficking and structure of RNA, and therefore making ribozymes work effectively in cells a significant challenge. This article addresses the problems of getting engineered ribozymes to effectively pair with and cleave targets in cells. The work described here illuminates methods for target-site selection on native mRNAs, methods for ribozyme expression, and strategies for obtaining a discrete intracellular localization of ribozymes.

L9 ANSWER 6 OF 15 MEDLINE DUPLICATE 2  
 ACCESSION NUMBER: 2001535732 MEDLINE  
 DOCUMENT NUMBER: 21467394 PubMed ID: 11582807  
 TITLE: Intracellular applications of ribozymes.  
 AUTHOR: \*\*\*Michienzi A\*\*\* ; Rossi J J  
 CORPORATE SOURCE: Molecular Biology Department, Beckman Research Institute of the City of Hope, Duarte, California 91010, USA.  
 CONTRACT NUMBER: AI 29329 (NIAID)  
 AI 42552 (NIAID)  
 AI 46030 (NIAID)  
 SOURCE: METHODS IN ENZYMOLOGY, (2001) 341 581-96.  
 Journal code: 0212271. ISSN: 0076-6879.  
 PUB. COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 200203  
 ENTRY DATE: Entered STN: 20011004  
 Last Updated on STN: 20020308  
 Entered Medline: 20020307

L9 ANSWER 7 OF 15 MEDLINE DUPLICATE 3  
 ACCESSION NUMBER: 2000422579 MEDLINE  
 DOCUMENT NUMBER: 20381318 PubMed ID: 10922055  
 TITLE: Ribozyme-mediated inhibition of HIV 1 suggests nucleolar trafficking of HIV-1 RNA.  
 AUTHOR: \*\*\*Michienzi A\*\*\* ; Cagnon L; Bahner I; Rossi J J  
 CORPORATE SOURCE: Department of Molecular Biology, Beckman Research Institute of the City of Hope, and Graduate School of Biological Sciences, City of Hope, Duarte, CA 91010-3011, USA.  
 CONTRACT NUMBER: 5 F32 GM18898-02 (NIGMS)  
 AI29329 (NIAID)  
 AI42552 (NIAID)  
 SOURCE: PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, (2000 Aug 1) 97 (16) 8955-60.  
 Journal code: 7505876. ISSN: 0027-8424.  
 PUB. COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals; AIDS  
 ENTRY MONTH: 200009  
 ENTRY DATE: Entered STN: 20000915  
 Last Updated on STN: 20000915  
 Entered Medline: 20000905

AB The HIV regulatory proteins Tat and Rev have a nucleolar localization property in human cells. However, no functional role has been attributed to this localization. Recently it has been demonstrated that expression of Rev induces nucleolar relocation of some protein factors involved in Rev export. Because the function of Rev is to bind HIV RNA and facilitate transport of singly spliced and unspliced RNA to the cytoplasm, it is likely that the nucleolus plays a critical role in HIV-1 RNA export. As a test for trafficking of HIV-1 RNAs into the nucleolus, a hammerhead ribozyme that specifically cleaves HIV-1 RNA was inserted into the body of the U16 small nucleolar RNA, resulting in accumulation of the ribozyme within the nucleoli of human cells. HeLa CD4(+) and T cells expressing this nucleolar localized ribozyme exhibit dramatically suppressed HIV-1 replication. The results presented here suggest a trafficking of HIV-1 RNA through the nucleoli of human cells, thus posing a different paradigm for lentiviral RNA processing.

L9 ANSWER 8 OF 15 MEDLINE DUPLICATE 4  
 ACCESSION NUMBER: 1999373046 MEDLINE  
 DOCUMENT NUMBER: 99373046 PubMed ID: 10445874  
 TITLE: The Rev protein is able to transport to the cytoplasm small nucleolar RNAs containing a Rev binding element.  
 AUTHOR: Buonomo S B; \*\*\*Michienzi A\*\*\* ; De Angelis F G; Bozzoni

CORPORATE SOURCE: Istituto Pasteur, Fondazione Cenci-Bolognetti, Dipartimento di Genetica e Biologia Molecolare, Universita di Roma La Sapienza, Italy.  
SOURCE: RNA, (1999 Aug) 5 (8) 993-1002.  
Journal code: 9509184. ISSN: 1355-8382.  
PUB. COUNTRY: United States  
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals; AIDS  
ENTRY MONTH: 199908  
ENTRY DATE: Entered STN: 19990913  
Last Updated on STN: 19990913  
Entered Medline: 19990830

AB Small nucleolar RNAs (snRNAs) were utilized to express Rev-binding sequences inside the nucleolus and to test whether they are substrates for Rev binding and transport. We show that U16 snRNA containing the minimal binding site for Rev stably accumulates inside the nucleolus maintaining the interaction with the basic C/D snRNA-specific factors. Upon Rev expression, the chimeric RNA is exported to the cytoplasm, where it remains bound to Rev in a particle devoid of snRNP-specific factors. These data indicate that Rev can elicit the functions of RNA binding and transport inside the nucleolus.

L9 ANSWER 9 OF 15 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 2000:108061 CAPLUS

DOCUMENT NUMBER: 132:248431

TITLE: A chimeric nucleolar Rev decoy inhibits the HIV replication

AUTHOR(S): \*\*\*Michienzi, A.\*\*\* ; Cagnon, L.; Bozzoni, I.; Rossi, J. J.

CORPORATE SOURCE: Department of Molecular Biology, Beckman Research

SOURCE: Institute of the City of Hope, Duarte, CA, 91010, USA  
Nucleic Acids Symposium Series (1999), 41(Symposium on RNA Biology III: RNA, Tool & Target), 211-214  
CODEN: NACSD8; ISSN: 0261-3166

PUBLISHER: Oxford University Press

DOCUMENT TYPE: Journal

LANGUAGE: English

AB In order to investigate the role of the nucleolus in Rev function, the authors constructed a nucleolar Rev decoy. They used the RBE (Rev binding element) sequence of the HIV genome as a decoy and U16snRNA as a vector to deliver the decoy into the nucleolar compartment (clone U16RBE). They then analyzed the U16RBE expression and anti-HIV activity in a human cell line. The data show that a nucleolar decoy is able to suppress the HIV replication, and suggest that the nucleolus can play a crit. role in Rev function.

REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 10 OF 15

MEDLINE

DUPLICATE 5

ACCESSION NUMBER: 1998211332 MEDLINE

DOCUMENT NUMBER: 98211332 PubMed ID: 9551610

TITLE: Inhibition of human immunodeficiency virus type 1 replication by nuclear chimeric anti-HIV ribozymes in a human T lymphoblastoid cell line.

AUTHOR: \*\*\*Michienzi A\*\*\* ; Conti L; Varano B; Prislei S; Gessani S; Bozzoni I

CORPORATE SOURCE: Istituto Pasteur, Fondazione Cenci-Bolognetti, Department of Genetics and Molecular Biology, University La Sapienza, Rome, Italy.

SOURCE: HUMAN GENE THERAPY, (1998 Mar 20) 9 (5) 621-8.  
Journal code: 9008950. ISSN: 1043-0342.

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English

FILE SEGMENT: Priority Journals; AIDS

ENTRY MONTH: 199805

ENTRY DATE: Entered STN: 19980611

Last Updated on STN: 19980611

Entered Medline: 19980529

AB Human immunodeficiency virus (HIV) infection represents one of the most challenging systems for gene therapy. Thanks to the extended knowledge of the molecular biology of the HIV life cycle, many different strategies have been developed including transdominant modifications of HIV proteins, RNA decoys, antisense RNA, ribozymes, and intracellular antibody fragments. In this paper, we have tested in a human T lymphoblastoid cell

- co-localize inside the nucleus with the Rev pre-mRNA before it is spliced and transported to the cytoplasm. This result was obtained by inserting the ribozyme in the spliceosomal U1 small nuclear RNA (snRNA) and in a derivative that has perfect complementarity with the 5' splice site of the Rev pre-mRNA. These ribozymes were tested in human T cell clones and were shown to be very efficient in inhibiting viral replication. Not only were the p24 levels in the culture medium drastically reduced but so were the intracellular HIV transcripts. Control disabled ribozymes enabled us to show the specificity of the ribozyme activity. Therefore, these constructs have potential utility for gene therapy of HIV-1 infection.

L9 ANSWER 11 OF 15 MEDLINE DUPLICATE 6  
 ACCESSION NUMBER: 97316827 MEDLINE  
 DOCUMENT NUMBER: 97316827 PubMed ID: 9174101  
 TITLE: Use of adenoviral VAI small RNA as a carrier for cytoplasmic delivery of ribozymes.  
 AUTHOR: Prislei S; Buonomo S B; \*\*\*Michienzi A\*\*\* ; Bozzoni I  
 CORPORATE SOURCE: Istituto Pasteur, Fondazione Cenci-Bolognetti, Dipartimento di Genetica e Biologia Molecolare, Universita La Sapienza, Roma, Italy.  
 SOURCE: RNA, (1997 Jun) 3 (6) 677-87.  
 Journal code: 9509184. ISSN: 1355-8382.  
 PUB. COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals; AIDS  
 ENTRY MONTH: 199706  
 ENTRY DATE: Entered STN: 19970709  
 Last Updated on STN: 19970709  
 Entered Medline: 19970624

AB The in vivo effectiveness of therapeutic RNAs, like antisense molecules and ribozymes, relies on several features: RNA molecules need to be expressed at high levels in the correct cellular compartment as stable and active molecules. The exploitation of "natural" small RNA coding genes as expressing cassettes gives high chances to fulfill these requirements. We have investigated the utilization of the adenoviral VAI RNA as a cytoplasmatic carrier for expressing ribozymes against HIV-1. The conserved 5' leader sequence of HIV was chosen as a target, because it is present in all the viral transcripts and is highly conserved. Hammerhead ribozymes were substituted to different portions of the VAI RNA and the resulting chimera were tested in the in vivo system of *Xenopus laevis* oocytes for their level of accumulation, cellular compartmentalization, and assembly in specific ribonucleoparticles containing the La antigen. Interesting differences in the activity of the different chimera were found in both in vitro cleavage assays and S100 extracts of injected oocytes where the catalytic activity of the ribozymes in the RNP context can be analyzed.

L9 ANSWER 12 OF 15 MEDLINE DUPLICATE 7  
 ACCESSION NUMBER: 96293503 MEDLINE  
 DOCUMENT NUMBER: 96293503 PubMed ID: 8692972  
 TITLE: U1 small nuclear RNA chimeric ribozymes with substrate specificity for the Rev pre-mRNA of human immunodeficiency virus.  
 AUTHOR: \*\*\*Michienzi A\*\*\* ; Prislei S; Bozzoni I  
 CORPORATE SOURCE: Istituto Pasteur Fondazione Cenci-Bolognetti, Rome, Italy.  
 SOURCE: PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, (1996 Jul 9) 93 (14) 7219-24.  
 Journal code: 7505876. ISSN: 0027-8424.  
 PUB. COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals; AIDS  
 ENTRY MONTH: 199608  
 ENTRY DATE: Entered STN: 19960911  
 Last Updated on STN: 19970203  
 Entered Medline: 19960829

AB The in vivo effectiveness of ribozymes strongly depends on the correct choice of the vector molecule. High levels of expression, stability, active conformation, and correct cellular localization are the most important features for a ribozyme vector. We have exploited the utilization of the U1 small nuclear RNA (snRNA) as a vector for specifically targeting a ribozyme into the nucleus. The Rev pre-mRNA of human immunodeficiency virus type 1 was chosen as target for testing the activity of the U1-ribozyme. The catalytic core of the hammerhead motif, plus the recognition sequences, substituted the stem-loop III of the U1

- vitro. In addition, in the in vivo system of *Xenopus laevis* oocytes, the U1-chimeric ribozyme accumulates in large amounts in the nucleus and produces a considerable reduction of Rev pre-mRNA levels. The Rev-specific ribozyme was also inserted in a derivative of the U1 snRNA mutated in the region of pairing with the 5' splice site, such as to match it with the suboptimal splice junction of the Rev precursor. This construct shows more efficient reduction of Rev pre-mRNA in vivo than the wild-type U1 vector.

L9 ANSWER 13 OF 15 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

ACCESSION NUMBER: 1994:281947 BIOSIS  
DOCUMENT NUMBER: PREV199497294947  
TITLE: Two novel small nucleolar RNA U16 and U18 are encoded inside introns of the same ribosomal protein gene.  
AUTHOR(S): Fragapane, Paola; Prislei, Silvia; \*\*\*Michienzi,\*\*\*  
\*\*\* Alessandro\*\*\* ; Presutti, Carlo; Caffarelli, Elisa; Bozzoni, Irene  
CORPORATE SOURCE: Centro Acidi Nucleici C.N.R., Dip. Genetica e Biol. Molecolare, Univ. "La Sapienza", Rome Italy  
SOURCE: Journal of Cellular Biochemistry Supplement, (1994) Vol. 0, No. 18C, pp. 113.  
Meeting Info.: Keystone Symposium on the Eukaryotic Nucleus  
Tamarro, Colorado, USA February 13-20, 1994  
ISSN: 0733-1959.  
DOCUMENT TYPE: Conference  
LANGUAGE: English

L9 ANSWER 14 OF 15 MEDLINE DUPLICATE 8

ACCESSION NUMBER: 94119676 MEDLINE  
DOCUMENT NUMBER: 94119676 PubMed ID: 7507233  
TITLE: Two different snRNAs are encoded in introns of amphibian and human L1 ribosomal protein genes.  
AUTHOR: Prislei S; \*\*\*Michienzi A\*\*\* ; Presutti C; Fragapane P; Bozzoni I  
CORPORATE SOURCE: Istituto Pasteur Fondazione Cenci-Bolognetti, Rome, Italy.  
SOURCE: NUCLEIC ACIDS RESEARCH, (1993 Dec 25) 21 (25) 5824-30.  
Journal code: 0411011. ISSN: 0305-1048.  
PUB. COUNTRY: ENGLAND: United Kingdom  
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
OTHER SOURCE: GENBANK-X75486; GENBANK-X75487; GENBANK-X75488  
ENTRY MONTH: 199402  
ENTRY DATE: Entered STN: 19940312  
Last updated on STN: 19960129  
Entered Medline: 19940222

AB We previously reported that the third intron of the *X.laevis* L1 ribosomal protein gene encodes for a snRNA called U16. Here we show that four different introns of the same gene contain another previously uncharacterized snRNA (U18) which is associated with fibrillarin in the nucleolus and which originates by processing of the pre-mRNA. The pathway of U18 RNA release from the pre-mRNA is the same as the one described for U16: primary endonucleolytic cleavages upstream and downstream of the U18 coding region produce a pre-U18 RNA which is subsequently trimmed to the mature form. Both the gene organization and processing of U18 are conserved in the corresponding genes of *X.tropicalis* and *H.sapiens*. The L1 gene thus has a composite structure, highly conserved in evolution, in which sequences coding for a ribosomal protein are intermingled with sequences coding for two different snRNAs. The nucleolar localization of these different components suggests some common function on ribosome biosynthesis.

L9 ANSWER 15 OF 15 MEDLINE DUPLICATE 9

ACCESSION NUMBER: 93327782 MEDLINE  
DOCUMENT NUMBER: 93327782 PubMed ID: 8335006  
TITLE: A novel small nucleolar RNA (U16) is encoded inside a ribosomal protein intron and originates by processing of the pre-mRNA.  
AUTHOR: Fragapane P; Prislei S; \*\*\*Michienzi A\*\*\* ; Caffarelli E; Bozzoni I  
CORPORATE SOURCE: Centro Acidi Nucleici of CNR, Universita La Sapienza, Roma, Italy.  
SOURCE: EMBO JOURNAL, (1993 Jul) 12 (7) 2921-8.  
Journal code: 8208664. ISSN: 0261-4189.  
PUB. COUNTRY: ENGLAND: United Kingdom  
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English



OTHER SOURCE: GENBANK-X72205  
ENTRY MONTH: 199308  
ENTRY DATE: Entered STN: 19930903  
Last Updated on STN: 19930903  
Entered Medline: 19930820

AB We report that the third intron of the L1 ribosomal protein gene of *Xenopus laevis* encodes a previously uncharacterized small nucleolar RNA that we called U16. This snRNA is not independently transcribed; instead it originates by processing of the pre-mRNA in which it is contained. Its sequence, localization and biosynthesis are phylogenetically conserved; in the corresponding intron of the human L1 ribosomal protein gene a highly homologous region is found which can be released from the pre-mRNA by a mechanism similar to that described for the amphibian U16 RNA. The presence of a snRNA inside an intron of the L1 ribosomal protein gene and the phylogenetic conservation of this gene arrangement suggest an important regulatory/functional link between these two components.

=> s l9 and tar  
L10 1 L9 AND TAR

=> d l10 ibib abs tot

L10 ANSWER 1 OF 1 MEDLINE  
ACCESSION NUMBER: 2002647867 MEDLINE  
DOCUMENT NUMBER: 22294961 PubMed ID: 12376617  
TITLE: A nucleolar \*\*\*TAR\*\*\* decoy inhibitor of HIV-1 replication.  
AUTHOR: \*\*\*Michienzi Alessandro\*\*\* ; Li Shirley; Zaia John A; Rossi John J  
CORPORATE SOURCE: Divisions of Molecular Biology and Virology, Beckman Research Institute of the City of Hope, 1450 East Duarte Road, Duarte, CA 91010-3011, USA.  
CONTRACT NUMBER: AI29329 (NIAID)  
SOURCE: AI46030 (NIAID) PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, (2002 Oct 29) 99 (22) 14047-52. Journal code: 7505876. ISSN: 0027-8424.  
PUB. COUNTRY: United States  
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 200212  
ENTRY DATE: Entered STN: 20021031  
Last Updated on STN: 20021217  
Entered Medline: 20021209

AB Tat is a critical regulatory factor in HIV-1 gene expression. It mediates the transactivation of transcription from the HIV-1 LTR by binding to the transactivation response ( \*\*\*TAR\*\*\* ) element in a complex with cyclin T1. Because of its critical and early role in HIV gene expression, Tat and its interaction with the \*\*\*TAR\*\*\* element constitute important therapeutic targets for the treatment of HIV-1 infection. Based on the known nucleolar localization properties of Tat, we constructed a chimeric small nucleolar RNA- \*\*\*TAR\*\*\* decoy that localizes to the nucleoli of human cells and colocalizes in the nucleolus with a Tat-enhanced GFP fusion protein. When the chimeric RNA was stably expressed in human T lymphoblastoid CEM cells it potentially inhibited HIV-1 replication. These results demonstrate that the nucleolar trafficking of Tat is critical for HIV-1 replication and suggests a role for the nucleolus in HIV-1 viral replication.

=> s tar and nucleoli  
L11 38 TAR AND NUCLEOLI

=> dup rem l11  
PROCESSING COMPLETED FOR L11  
L12 21 DUP REM L11 (17 DUPLICATES REMOVED)

=> s l12 not l10  
L13 20 L12 NOT L10

=> d l13 ibib abs tot

L13 ANSWER 1 OF 20 MEDLINE  
ACCESSION NUMBER: 97375656 MEDLINE

TITLE: . Xlrpba, a double-stranded RNA-binding protein associated with ribosomes and heterogeneous nuclear RNPs.  
 AUTHOR: Eckmann C R; Jantsch M F  
 CORPORATE SOURCE: Department of Cytology and Genetics, Institute of Botany, University of Vienna, A-1030 Vienna, Austria.  
 SOURCE: JOURNAL OF CELL BIOLOGY, (1997 Jul 28) 138 (2) 239-53.  
 Journal code: 0375356. ISSN: 0021-9525.  
 PUB. COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals; AIDS  
 ENTRY MONTH: 199708  
 ENTRY DATE: Entered STN: 19970908  
 Last Updated on STN: 19970908  
 Entered Medline: 19970825

AB we have cloned and characterized Xlrpba, a double-stranded RNA-binding protein from *Xenopus laevis*. Xlrpba is a protein of 33 kD and contains three tandemly arranged, double-stranded RNA-binding domains (dsRBDs) that bind exclusively to double-stranded RNA in vitro, but fail to bind either single-stranded RNA or DNA. Sequence data and the overall organization of the protein suggest that Xlrpba is the *Xenopus* homologue of human \*\*\*TAR\*\*\* -RNA binding protein (TRBP), a protein isolated by its ability to bind to human immunodeficiency virus (HIV) \*\*\*TAR\*\*\* -RNA. In transfection assays, TRBP has also been shown to inhibit the interferon-induced protein kinase PKR possibly by direct physical interaction. To determine the function of Xlrpba and its human homologue we studied the expression and intracellular distribution of the two proteins. Xlrpba is ubiquitously expressed with marked quantitative differences amongst all tissues. Xlrpba and human TRBP can be detected in the cytoplasm and nucleus by immunofluorescence staining and western blotting. Sedimentation gradient analyses and immunoprecipitation experiments suggest an association of cytoplasmic Xlrpba with ribosomes. In contrast, a control construct containing two dsRBDs fails to associate with ribosomes in microinjected *Xenopus* oocytes. Nuclear staining of *Xenopus* lampbrush chromosome preparations showed the association of the protein with \*\*\*nucleoli\*\*\*, again indicating an association of the protein with ribosomal RNAs. Additionally, Xlrpba could be located on lampbrush chromosomes and in snurposomes. Immunoprecipitations of nuclear extracts demonstrated the presence of the protein in heterogeneous nuclear (hn) RNP particles, but not in small nuclear RNPs, explaining the chromosomal localization of the protein. It thus appears that Xlrpba is a general double-stranded RNA-binding protein which is associated with the majority of cellular RNAs, ribosomal RNAs, and hnRNAs either alone or as part of an hnRNP complex.

L13 ANSWER 2 OF 20 MEDLINE  
 ACCESSION NUMBER: 97360029 MEDLINE  
 DOCUMENT NUMBER: 97360029 PubMed ID: 9217057  
 TITLE: Bovine immunodeficiency virus tat gene: cloning of two distinct cDNAs and identification, characterization, and immunolocalization of the tat gene products.  
 AUTHOR: Fong S E; Greenwood J D; Williamson J C; Derse D; Pallansch L A; Copeland T; Rasmussen L; Mentzer A; Nagashima K; Tobin G; Gonda M A  
 CORPORATE SOURCE: Laboratory of Cell and Molecular Structure, SAIC Frederick, NCI-Frederick Cancer Research and Development Center, Maryland 21702-1201, USA.. Fong@mail.ncifcrf.gov  
 SOURCE: VIROLOGY, (1997 Jul 7) 233 (2) 339-57.  
 Journal code: 0110674. ISSN: 0042-6822.  
 PUB. COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals; AIDS  
 ENTRY MONTH: 199708  
 ENTRY DATE: Entered STN: 19970825  
 Last Updated on STN: 19970825  
 Entered Medline: 19970813

AB cDNAs encoding the bovine immunodeficiency virus (BIV) transactivator gene (tat) were cloned from virally infected cells and characterized. BIV expresses two distinct tat mRNAs composed of three exons that are derived by alternative splicing. The BIV tat mRNA splice variants encode Tat proteins of 103 (Tat103) and 108 (Tat108) amino acids. The Tat103 coding region is specified only by exon 2, while that of Tat108 is specified by a truncated exon 2 and the first 30 nt of exon 3. Thus, the first 98 amino acids of each Tat are identical, and have amino terminal, cysteine-rich, conserved core, basic, and carboxyl-terminal domains similar to Tats

14-kDa phosphorylated Tat protein identical in size to recombinant Tat expressed in bacteria. BIV Tat was shown to localize exclusively in the \*\*\*nucleoli\*\*\* of virally infected and Tat-expressing cells. Reporter gene assays indicated that Tat103 and Tat108 can strongly transactivate the BIV long terminal repeat (LTR) in virally permissive canine Cf2Th and nonpermissive HeLa and mouse NIH 3T3 cells, but not in permissive lapine EREp cells. However, an intact BIV tat gene is required for viral replication in both Cf2Th and EREp cells. Strong LTR activation by BIV Tat requires a \*\*\*TAR\*\*\* (transactivation responsive) element delimited by viral nt +1 to +31 and the Tat basic domain. BIV Tat strongly cross-transactivates the HIV-1 LTR in a \*\*\*TAR\*\*\* -dependent manner in Cf2Th, but not in EREp, HeLa, or NIH 3T3 cells. In contrast, strong, \*\*\*TAR\*\*\* -dependent cross-transactivation of the BIV LTR by HIV-1 Tat could not be demonstrated in any of these cell types. In Cf2Th cells Tat108 effects a moderately stronger transactivation of the BIV LTR than Tat103, indicative of a functional difference in BIV Tat proteins encoded by the mRNA splice variants. The present studies demonstrate that BIV Tat parallels the primate lentiviral Tats in structure and biochemistry but is not interchangeable with the latter.

L13 ANSWER 3 OF 20 MEDLINE

ACCESSION NUMBER: 91175688 MEDLINE  
DOCUMENT NUMBER: 91175688 PubMed ID: 2078551  
TITLE: Regulation of expression of human immunodeficiency virus.  
AUTHOR: Pavlakakis G N; Felber B K  
CORPORATE SOURCE: National Cancer Institute, Frederick Cancer Research Facility, BRI-Basic Research Program, MD 21701-1013.  
CONTRACT NUMBER: N01-CO-74101 (NCI)  
SOURCE: NEW BIOLOGIST, (1990 Jan) 2 (1) 20-31. Ref: 141  
Journal code: 9000976. ISSN: 1043-4674.  
PUB. COUNTRY: United States  
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
General Review; (REVIEW)  
(REVIEW, ACADEMIC)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals; AIDS  
ENTRY MONTH: 199105  
ENTRY DATE: Entered STN: 19910519  
Last Updated on STN: 19970203  
Entered Medline: 19910501

AB Human immunodeficiency virus (HIV), the causative agent of acquired immune deficiency syndrome (AIDS), encodes regulatory factors necessary for its expression. The study of these factors clearly indicates the importance and complexity of post-transcriptional processes in regulating gene expression. Two regulatory proteins, Tat and Rev, are necessary for viral replication. These small nuclear proteins accumulate primarily in the \*\*\*nucleoli\*\*\* and act on HIV via sequence-specific elements found on the viral RNA. Both the Tat responsive element ( \*\*\*TAR\*\*\* ) and the Rev responsive element (RRE) map within regions of strong RNA secondary structure. While all viral mRNAs contain \*\*\*TAR\*\*\*, only the mRNAs producing HIV structural proteins contain RRE. Tat increases the levels of all viral mRNAs. Its function is complex and involves transcriptional and possibly post-transcriptional steps. Rev promotes the transport of viral mRNAs containing RRE from the nucleus to the cytoplasm and increases the half-life of these viral mRNAs. A third factor, Nef, is a cytoplasmic myristylated protein that has been proposed to down-regulate virus expression. These factors are integrated in a feedback regulatory network that dictates the balanced expression of viral components. The study of HIV expression in human cells will advance our understanding of the mechanisms related to the pathogenicity of HIV. This knowledge may also lead to novel diagnostic and therapeutic approaches to combat AIDS.

L13 ANSWER 4 OF 20 MEDLINE

ACCESSION NUMBER: 80094875 MEDLINE  
DOCUMENT NUMBER: 80094875 PubMed ID: 521529  
TITLE: Ultrastructural and cell surface changes of human psoriatic skin following Goeckerman therapy.  
AUTHOR: Lupulescu A P; Chadwick J M; Downham T F 2nd  
SOURCE: JOURNAL OF CUTANEOUS PATHOLOGY, (1979 Oct) 6 (5) 347-63.  
Journal code: 0425124. ISSN: 0303-6987.  
PUB. COUNTRY: Denmark  
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 198003  
ENTRY DATE: Entered STN: 19900315

AB Ultrastructural and cell surface studies of skin in psoriatic patients prior to and after Goeckerman therapy (crude coal **\*\*\*tar\*\*\*** and UVB-light) have demonstrated significant cellular changes following this treatment: hyperactivity of melanocytes with melanosome polymorphism, increase of desmosomes, tonofilaments, keratohyaline granules, a decrease in mitochondria, keratinosomes, polysomes, dark cells and a reduction in size of nuclei and **\*\*\*nucleoli\*\*\***. The enlargement of intercellular spaces and the redundancy of basement membrane were also reduced. Langerhans cells were moderately decreased and exhibited a normal ultrastructural pattern. No significant changes in cutaneous nerve distribution or morphology were observed in these cases. Scanning electron microscopy following treatment revealed a regular surface and orientation of corneocytes, with flattened surfaces; and a reduction of their ridges on the surfaces, as well as of the intercellular spaces and red blood cells. These findings indicate that Goeckerman therapy restored the ultrastructural and cell surface pattern in the psoriatic skin by inducing orthokeratinogenesis, development of the tonofibrillar-desmosome system, and decrease in mitochondria, nuclei and **\*\*\*nucleoli\*\*\***.

L13 ANSWER 5 OF 20 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1984:485369 CAPLUS

DOCUMENT NUMBER: 101:85369

TITLE: The effects of **\*\*\*tar\*\*\*** dyes on RNA synthesis. Part 5. Configurational changes in rat liver nuclear chromatin caused by azo dyes

AUTHOR(S): Yoshimoto, M.; Yamaguchi, M.; Hatano, S.; Watanabe, T.

CORPORATE SOURCE: Fac. Agric., Kyushu Univ., Fukuoka, 812, Japan

SOURCE: Food and Chemical Toxicology (1984), 22(5), 3374-44

CODEN: FCTOD7; ISSN: 0278-6915

DOCUMENT TYPE: Journal

LANGUAGE: English

GI

/ structure 1 in file .gra /

AB The effects of azo dyes on the chromatin configuration of isolated rat liver nuclei were examd. by electron microscopy for elucidation of the mechanism of dye-stimulated in vitro RNA synthesis. Sunset Yellow FCF [2783-94-0] had no effect on the incorporation of [3H]UTP into RNA but amaranth (I) [915-67-3] and Ponceau 3R [3564-09-8] stimulated incorporation .apprx.2- and 3.5-fold, resp. Sunset Yellow FCF did not produce any major alteration in liver nuclei. Isolated liver nuclei treated with I or Ponceau 3R showed dissocn. of heterochromatin and the **\*\*\*nucleoli\*\*\*** lost their fine-granular structure acquiring a coarse-granular configuration. The dense structure of heterochromatin in liver nuclei treated with Ponceau 3R was completely dissocd. and appeared as a lace-like configuration. Thus, both Ponceau 3R and I stimulate in vitro RNA synthesis by causing the dissocn. of heterochromatin in isolated rat liver nuclei.

L13 ANSWER 6 OF 20 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1964:495149 CAPLUS

DOCUMENT NUMBER: 61:95149

ORIGINAL REFERENCE NO.: 61:16588c-e

TITLE: Content and state of nucleic acids in tissues undergoing inflammatory processes

AUTHOR(S): Varfolomeeva, Z. N.

SOURCE: Biol. Nukleinovogo Obmena u Rast., Akad. Nauk SSSR, Bashkirsk. Filial, Materialy 2-oi Nauchn. [Vtoroi] Konf., Ufa (1964), 1962, 111-14

DOCUMENT TYPE: Journal

LANGUAGE: Unavailable

AB A rise in nucleic acids is known to occur in recovering wound tissue of both plants and animals. In regenerating cells, labile ribonucleic acid predominates in the cytoplasm; in differentiated tissue, the stable form. Labile deoxyribonucleic acid (DNA) in the nucleus stains red in the pyronine-methyl green test; the stable, green. In rabbit pancreas subjected to a coal **\*\*\*tar\*\*\*** irritant, cells in direct contact necrosed, with disappearing nucleic acids, as did older ones at some distance, but younger ones passed through 3 stages of change: (1) regression or dedifferentiation, (2) proliferation, and (3) redifferentiation. In 1 the nucleic acids fell sharply in the epithelial cells. In 2 they rose above normal; the inflammatory zone was

\*\*\*nucleoli\*\*\* (red). In 3 a gradual return to normal occurred, with the nucleus still staining green and \*\*\*nucleoli\*\*\* absent.

L13 ANSWER 7 OF 20 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1956:21176 CAPLUS  
DOCUMENT NUMBER: 50:21176  
ORIGINAL REFERENCE NO.: 50:4373b-d  
TITLE: Early cytologic changes produced by carcinogens  
AUTHOR(S): Cooper, Norman S.  
CORPORATE SOURCE: New York Univ.-Bellvue Med. Center, New York, NY  
SOURCE: Bull. N.Y. Acad. Med. (1956), 32, 79-80  
DOCUMENT TYPE: Journal  
LANGUAGE: Unavailable

AB Enlargement of mouse epidermal cell \*\*\*nucleoli\*\*\* was used as a criterion of the carcinogenicity of a substance painted on the skin of the animal. U.S.P. white mineral oil caused enlargement, therefore only materials giving enlargements greater than this were considered pos. Cigaret-smoke condensate (50% in acetone) was pos.; 50% coal \*\*\*\*tar\*\*\* in acetone gave even greater enlargement. The greatest enlargement was given by 0.5% 3,4-benzopyrene in acetone. A 50% soln. of turpentine in acetone produced nucleolar enlargement only slightly less than that resulting from tobacco \*\*\*\*tar\*\*\*.

L13 ANSWER 8 OF 20 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1928:18650 CAPLUS  
DOCUMENT NUMBER: 22:18650  
ORIGINAL REFERENCE NO.: 22:2199b-e  
TITLE: Studies in the microchemistry of the cell. I. The chromatin content of normal and malignant cells as demonstrated by Feulgen's "Nuclearreaktion"  
AUTHOR(S): Ludford, R. J.  
SOURCE: Proc. Roy. Soc. (London) (1928), B102, 397-406  
DOCUMENT TYPE: Journal  
LANGUAGE: Unavailable

AB The following results were obtained by use of this reaction. In the rat and the mouse, the chromatin content of the nucleus does not increase during oogenesis, and no chromatin is extruded into the cytoplasm when the germinal vesicle breaks down to form the chromosomes. Chromatin is not present in either the oxyphilic or the basophilic \*\*\*nucleoli\*\*\* of the oocyte of the mollusk, Limnoea stagnalis. The sperm heads of all 3 species yield a faint reaction when the chromosomes are stretched; but the reaction becomes progressively more marked as condensation occurs in spermatogenesis. The chromosomes apparently contain other substances besides chromatin. Extrusion of chromatin does not occur in the epithelial cells of the epididymis. Shrinkage of the nuclei of gland cells after secretion (e. g., in the adrenal medulla after exposure to cold) is apparently due to the loss of some substance other than chromatin. No relationship is found between the chromatin content of a tumor cell nucleus and the rate of growth of the tumor. In a \*\*\*\*tar\*\*\* tumor and the surrounding skin, the chromatin content is apparently the same in the normal and the malignant cells. During cellular degeneration of tumors, their nuclei become shrunken and the chromatin runs together. In tumors, the chromosomes stain intensely, and are separate from the nucleolus during the prophase of mitosis. Nuclear extrusions, which are well marked in some tumors, consist of nucleolar material and not chromatin. Nucleolar extrusions are the chief source of keratohyalin during cornification. While the same amt. of chromatin may be present in both large and small nuclei, yet giant nuclei contain large masses of chromatin.

L13 ANSWER 9 OF 20 USPATFULL

ACCESSION NUMBER: 2002:279998 USPATFULL  
TITLE: Genetically engineered herpes virus for the treatment of cardiovascular disease  
INVENTOR(S): Schwartz, Lewis B., Hinsdale, IL, UNITED STATES  
Weichselbaum, Ralph R., Chicago, IL, UNITED STATES  
Roizman, Bernard, Chicago, IL, UNITED STATES

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 2002155432	A1	20021024
APPLICATION INFO.:	US 2001-995475	A1	20011128 (9)

	NUMBER	DATE
PRIORITY INFORMATION:	US 2000-253680P	20001128 (60)

FILE SEGMENT: APPLICATION  
LEGAL REPRESENTATIVE: MARSHALL, GERSTEIN & BORUN, 6300 SEARS TOWER, 233 SOUTH  
WACKER, CHICAGO, IL, 60606-6357  
NUMBER OF CLAIMS: 33  
EXEMPLARY CLAIM: 1  
LINE COUNT: 4203

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB The present invention provides methods of expressing a nucleic acid or producing a proteinaceous composition encoded by a nucleic acid in vascular and cardiovascular cells by administration of a herpesvirus vector. The present invention provides methods of producing a therapeutic benefit in vascular and cardiovascular tissue by administration of a herpesvirus vector. In additional aspects, the invention concerns combination therapies for vascular and cardiovascular diseases comprising administration of a herpesvirus vector and treatment with at least one addition pharmacological agent or surgical procedure.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L13 ANSWER 10 OF 20 USPATFULL

ACCESSION NUMBER: 2002:217408 USPATFULL  
TITLE: Oligomers that bind to ku protein  
INVENTOR(S): Dynan, William S., Augusta, GA, United States  
Yoo, Sunghan, Augusta, GA, United States  
PATENT ASSIGNEE(S): Medical College of Georgia Research Institute, Inc.,  
Augusta, GA, United States (U.S. corporation)

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 6441158	B1	20020827
APPLICATION INFO.:	US 1998-223139		19981230 (9)

	NUMBER	DATE
PRIORITY INFORMATION:	US 1997-70278P	19971231 (60)
DOCUMENT TYPE:	Utility	
FILE SEGMENT:	GRANTED	
PRIMARY EXAMINER:	Wang, Andrew	
LEGAL REPRESENTATIVE:	Kilpatrick Stockton LLP, Rothschild, Cynthia B., Calkins, Charles W.	
NUMBER OF CLAIMS:	8	
EXEMPLARY CLAIM:	1	
NUMBER OF DRAWINGS:	8 Drawing Figure(s); 5 Drawing Page(s)	
LINE COUNT:	2333	

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB Disclosed are oligomers that bind Ku protein. These oligomers are useful for inhibiting activation of DNA-PK, treating certain forms of autoimmune disease, detection and purification of Ku protein, and identification of proteins that interact with Ku protein. Preferably, the oligomers are composed of nucleotides, nucleotide analogs, or a combination. Most preferably, the oligomers are composed of ribonucleotides. Also disclosed is a method of inhibiting DNA repair, a method of identifying cellular proteins that interact with Ku protein, and a method of treating autoimmune disease in patients with anti-Ku antibodies. The disclosed oligomers can have several preferred features, either alone or in combination, in addition to Ku binding. One such feature, referred to herein as inhibition activity, is inhibition of DNA-PK kinase activity. Another preferred feature, referred to herein as aptamer motifs, is the presence of one or more of the base sequences GCUUCCCCANNAC, A(A/C)AUGA, and AACUUCGA. These sequences--referred to herein as aptamer motif 1, aptamer motif 2, and aptamer motif 3, respectively--are associated with Ku binding capability. Another preferred feature, referred to herein as aptamer structure, is the presence of a structure similar to the structure shown in FIG. 6A. This structure has the general formula 5'-A-B-C-D-C'-E-A'-3', where A, B, C, D, C', E, and A' are components of the oligomer. In this structure, A and A' interact to form a stem structure, C and C' interact to form a stem structure, B and E make up a bulge region, and D is either a bulge or a loop. FIG. 6A depicts component D as a loop. Each of these preferred features (inhibition activity, aptamer motif, and aptamer structure) can be used either alone or in combination with one or both of the other characteristics.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L13 ANSWER 11 OF 20 USPATFULL

TITLE: . HIV chemokines  
INVENTOR(S): Ludwig, Linda B., East Aurora, NY, United States  
Ambrus, Jr., Julian L., Buffalo, NY, United States  
Krawczyk, Kristie Anne, Gowanda, NY, United States  
PATENT ASSIGNEE(S): The Research Foundation of State University of New  
York, Amherst, NY, United States (U.S. corporation)

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 6392029	B1	20020521
APPLICATION INFO.:	US 1999-249542		19990212 (9)
RELATED APPLN. INFO.:	Continuation-in-part of Ser. No. US 1997-853703, filed on 9 May 1997, now patented, Pat. No. US 5919677		

	NUMBER	DATE
PRIORITY INFORMATION:	US 1998-74640P	19980213 (60)
DOCUMENT TYPE:	Utility	
FILE SEGMENT:	GRANTED	
PRIMARY EXAMINER:	Bui, Phuong T.	
LEGAL REPRESENTATIVE:	Hodgson Russ LLP	
NUMBER OF CLAIMS:	7	
EXEMPLARY CLAIM:	1	
NUMBER OF DRAWINGS:	19 Drawing Figure(s); 16 Drawing Page(s)	
LINE COUNT:	1868	

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB Disclosed is a gene comprising an open reading frame encoded on the plus strand of the pro-viral DNA, and located in the region of HIV-1 long terminal repeat. The gene encodes a protein that is related to, and has a structural motif resembling that of chemokine proteins. Depending upon the ribosomal frameshift, a plurality of proteins may be translated from the antisense RNA. The protein has similarity with chemokine SDF-1 and may play a role as a cofactor with gp120 in the binding to and entry of HIV to a target cell.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L13 ANSWER 12 OF 20 USPATFULL

ACCESSION NUMBER: 2001:202200 USPATFULL  
TITLE: Tat-derived transport polypeptides  
INVENTOR(S): Frankel, Alan, Tiburon, CA, United States  
Pabo, Carl, Newton, MA, United States  
Barsom, James G., Lexington, MA, United States  
Fawell, Stephen E., Winchester, MA, United States  
Pepinsky, R. Blake, Arlington, MA, United States  
PATENT ASSIGNEE(S): Whitehead Institute for Biomedical Research, Cambridge,  
MA, United States (U.S. corporation)  
Johns Hopkins Univ. School of Medicine, Baltimore, MD,  
United States (U.S. corporation)  
Biogen, Inc., Cambridge, MA, United States (U.S.  
corporation)

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 6316003	B1	20011113
APPLICATION INFO.:	US 1994-235403		19940428 (8)
RELATED APPLN. INFO.:	Continuation-in-part of Ser. No. US 1993-158015, filed on 24 Nov 1993, now abandoned Continuation of Ser. No. US 1991-636662, filed on 2 Jan 1991, now abandoned Continuation-in-part of Ser. No. US 1989-454450, filed on 21 Dec 1989, now abandoned, said Ser. No. US 636662 And Ser. No. WO 1993-US7833, filed on 19 Aug 1993, said Ser. No. US 636662 And Ser. No. US 1992-934375, filed on 21 Aug 1992, now abandoned		
DOCUMENT TYPE:	Utility		
FILE SEGMENT:	GRANTED		
PRIMARY EXAMINER:	MacMillan, Keith D.		
ASSISTANT EXAMINER:	Wessendorf, T. D.		
NUMBER OF CLAIMS:	11		
EXEMPLARY CLAIM:	1		
NUMBER OF DRAWINGS:	21 Drawing Figure(s); 16 Drawing Page(s)		
LINE COUNT:	3392		

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB This invention relates to delivery of biologically active cargo molecules, such as polypeptides and nucleic acids, into the cytoplasm and nuclei of cells in vitro and in vivo. Intracellular delivery of

of novel transport polypeptides which include HIV tat protein or one or more portions thereof, and which are covalently attached to cargo molecules. The transport polypeptides in preferred embodiments of this invention are characterized by the presence of the tat basic region (amino acids 49-57), the absence of the tat cysteine-rich region (amino acids 22-36) and the absence of the tat exon 2-encoded carboxy-terminal domain (amino acids 73-86) of the naturally-occurring tat protein. By virtue of the absence of the cysteine-rich region, the preferred transport polypeptides of this invention solve the potential problems of spurious trans-activation and disulfide aggregation. The reduced size of the preferred transport polypeptides of this invention also minimizes interference with the biological activity of the cargo molecule.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L13 ANSWER 13 OF 20 USPATFULL

ACCESSION NUMBER: 1998:108434 USPATFULL  
TITLE: Tat-derived transport polypeptides and fusion proteins  
INVENTOR(S): Frankel, Alan, Tiburon, CA, United States  
Pabo, Carl, Newton, MA, United States  
Barsoum, James G., Lexington, MA, United States  
Fawell, Stephen E., Winchester, MA, United States  
Pepinsky, R. Blake, Arlington, MA, United States  
PATENT ASSIGNEE(S): Biogen, Inc., Cambridge, MA, United States (U.S. corporation)

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 5804604		19980908
APPLICATION INFO.:	US 1995-450236		19950525 (8)
RELATED APPLN. INFO.:	Continuation of Ser. No. US 1994-235403, filed on 28 Apr 1994 which is a continuation-in-part of Ser. No. US 1993-158015, filed on 24 Nov 1993, now abandoned which is a continuation of Ser. No. US 1991-636662, filed on 2 Jan 1991, now abandoned which is a continuation-in-part of Ser. No. US 1989-454450, filed on 21 Dec 1989, now abandoned		
DOCUMENT TYPE:	Utility		
FILE SEGMENT:	Granted		
PRIMARY EXAMINER:	Elliott, George C.		
ASSISTANT EXAMINER:	Larson, Thomas G.		
LEGAL REPRESENTATIVE:	Biogen, Inc., Kaplan, Warren A.		
NUMBER OF CLAIMS:	14		
EXEMPLARY CLAIM:	1		
NUMBER OF DRAWINGS:	26 Drawing Figure(s); 21 Drawing Page(s)		
LINE COUNT:	4456		

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB This invention relates to delivery of biologically active cargo molecules, such as polypeptides and nucleic acids, into the cytoplasm and nuclei of cells in vitro and in vivo. Intracellular delivery of cargo molecules according to this invention is accomplished by the use of novel transport polypeptides which comprise HIV tat protein or one or more portions thereof, and which are covalently attached to cargo molecules. The transport polypeptides in preferred embodiments of this invention are characterized by the presence of the tat basic region (amino acids 49-57), the absence of the tat cysteine-rich region (amino acids 22-36) and the absence of the tat exon 2-encoded carboxy-terminal domain (amino acids 73-86) of the naturally-occurring tat protein. By virtue of the absence of the cysteine-rich region, the preferred transport polypeptides of this invention solve the potential problems of spurious trans-activation and disulfide aggregation. The reduced size of the preferred transport polypeptides of this invention also minimizes interference with the biological activity of the cargo molecule.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L13 ANSWER 14 OF 20 USPATFULL

ACCESSION NUMBER: 1998:48556 USPATFULL  
TITLE: Tat-derived transport polypeptide conjugates  
INVENTOR(S): Frankel, Alan, 21 Marinero Cir., #206, Tiburon, CA, United States 94920  
Pabo, Carl, 18 Weldon Rd., Newton, MA, United States 02158  
Barsoum, James G., 9 Marlboro Rd., Lexington, MA, United States 02173  
Fawell, Stephen E., One Black Horse Ter., Winchester,



Pepinsky, R. Blake, 30 Falmouth Rd., Arlington, MA,  
United States 02174

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 5747641		19980505
APPLICATION INFO.:	US 1995-451233		19950525 (8)
RELATED APPLN. INFO.:	Continuation of Ser. No. US 1994-235403, filed on 28 Apr 1994 which is a continuation-in-part of Ser. No. US 1993-158015, filed on 24 Nov 1993, now abandoned which is a continuation of Ser. No. US 1991-636662, filed on 2 Jan 1991, now abandoned which is a continuation-in-part of Ser. No. US 1989-454450, filed on 21 Dec 1989, now abandoned And a continuation-in-part of Ser. No. US -235403 which is a continuation-in-part of Ser. No. US 1992-934375, filed on 12 Aug 1992, now abandoned		
DOCUMENT TYPE:	Utility		
FILE SEGMENT:	Granted		
PRIMARY EXAMINER:	Elliott, George C.		
ASSISTANT EXAMINER:	Larson, Thomas G.		
LEGAL REPRESENTATIVE:	Fish & Neave, Haley, Jr., James F., Kanter, Madge R.		
NUMBER OF CLAIMS:	5		
EXEMPLARY CLAIM:	1		
NUMBER OF DRAWINGS:	21 Drawing Figure(s); 16 Drawing Page(s)		
LINE COUNT:	3850		
CAS INDEXING IS AVAILABLE FOR THIS PATENT.			

AB This invention relates to delivery of biologically active cargo molecules, such as polypeptides and nucleic acids, into the cytoplasm and nuclei of cells in vitro and in vivo. Intracellular delivery of cargo molecules according to this invention is accomplished by the use of novel transport polypeptides which comprise HIV tat protein or one or more portions thereof, and which are covalently attached to cargo molecules. The transport polypeptides in preferred embodiments of this invention are characterized by the presence of the tat basic region (amino acids 49-57), the absence of the tat cysteine-rich region (amino acids 22-36) and the absence of the tat exon 2-encoded carboxy-terminal domain (amino acids 73-86) of the naturally-occurring tat protein. By virtue of the absence of the cysteine-rich region, the preferred transport polypeptides of this invention solve the potential problems of spurious trans-activation and disulfide aggregation. The reduced size of the preferred transport polypeptides of this invention also minimizes interference with the biological activity of the cargo molecule.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L13 ANSWER 15 OF 20 USPATFULL  
ACCESSION NUMBER: 97:106804 USPATFULL  
TITLE: Cellular protein TDP-43 and regulation of HIV-1 gene expression  
INVENTOR(S): Gaynor, Richard B., Dallas, TX, United States  
Ou, S.-H. Ignatius, Dallas, TX, United States  
Wu, Foon Kin, Carrollton, TX, United States  
PATENT ASSIGNEE(S): Board of Regents, The University of Texas System,  
Austin, TX, United States (U.S. corporation)

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 5688511		19971118
APPLICATION INFO.:	US 1994-343682		19941122 (8)
RELATED APPLN. INFO.:	Continuation-in-part of Ser. No. US 1991-788266, filed on 5 Nov 1991, now patented, Pat. No. US 5350835 And Ser. No. US 1994-239047, filed on 6 May 1994, now abandoned		
DOCUMENT TYPE:	Utility		
FILE SEGMENT:	Granted		
PRIMARY EXAMINER:	Nucker, Christine M.		
ASSISTANT EXAMINER:	Stucker, Jeffrey		
LEGAL REPRESENTATIVE:	Arnold, White & Durkee		
NUMBER OF CLAIMS:	20		
EXEMPLARY CLAIM:	1		
NUMBER OF DRAWINGS:	2 Drawing Figure(s); 2 Drawing Page(s)		
LINE COUNT:	2863		
CAS INDEXING IS AVAILABLE FOR THIS PATENT.			

AB Compositions including a polypeptide or nucleic acid sequence encoding a polypeptide that binds \*\*\*TAR\*\*\* DNA (particularly the region -18 to

.(particularly the region +1 to +80 of the \*\*\*TAR\*\*\* RNA) are disclosed. The cellular binding protein TDP-43 including the polypeptide has an estimated molecular weight of between about 40 kD and 46 kD as determined by SDS polyacrylamide gel electrophoresis. Fusion proteins that include the entire cellular binding protein TDP-43 or fragments thereof are also described. The cellular binding protein, peptide fragments and nucleic acid sequences encoding them, repress HIV gene expression. Methods for preparing the cellular binding protein from cells as recombinant proteins with recombinant host cells are also disclosed. Antibodies to the TDP-43 cellular binding protein are also described. The isolated nucleic acid sequences of the protein and its fragments are described in the construction of retroviral vectors. Methods for using the cellular binding protein, polypeptides thereof, and nucleic acid sequences encoding the protein and its fragments in the preparation of pharmaceutical preparations and in methods for repressing HIV gene expression are also disclosed.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L13 ANSWER 16 OF 20 USPATFULL

ACCESSION NUMBER: 97:91635 USPATFULL

TITLE: Fusion protein comprising tat-derived transport moiety

INVENTOR(S): Frankel, Alan, 21 Marinero Cir. #206, Tiburon, CA, United States 94920

Pabo, Carl, 18 Weldon Rd., Newton, MA, United States 02158

Barsoum, James G., 9 Marlboro Rd., Lexington, MA, United States 02173

Fawell, Stephen E., One Black Horse Ter., Winchester, MA, United States 01890

Pepinsky, R. Blake, 30 Falmouth Rd., Arlington, MA, United States 02174

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 5674980		19971007
APPLICATION INFO.:	US 1995-450098		19950525 (8)
RELATED APPLN. INFO.:	Continuation of Ser. No. US 1994-235403, filed on 28 Apr 1994, now abandoned which is a continuation-in-part of Ser. No. US 1993-158015, filed on 24 Nov 1993, now abandoned which is a continuation of Ser. No. US 1991-636662, filed on 2 Jan 1991, now abandoned which is a continuation-in-part of Ser. No. US 1989-454450, filed on 21 Dec 1989, now abandoned, said Ser. No. US -235403 which is a continuation-in-part of Ser. No. US 1992-934375, filed on 21 Aug 1992, now abandoned		
DOCUMENT TYPE:	Utility		
FILE SEGMENT:	Granted		
PRIMARY EXAMINER:	Elliott, George C.		
ASSISTANT EXAMINER:	Larson, Thomas G.		
LEGAL REPRESENTATIVE:	Fish & Neave, Haley, Jr., James F., Kanter, Madge R.		
NUMBER OF CLAIMS:	10		
EXEMPLARY CLAIM:	1		
NUMBER OF DRAWINGS:	21 Drawing Figure(s); 16 Drawing Page(s)		
LINE COUNT:	3855		

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB This invention relates to delivery of biologically active cargo molecules, such as polypeptides and nucleic acids, into the cytoplasm and nuclei of cells in vitro and in vivo. Intracellular delivery of cargo molecules according to this invention is accomplished by the use of novel transport polypeptides which comprise HIV tat protein or one or more portions thereof, and which are covalently attached to cargo molecules. The transport polypeptides in preferred embodiments of this invention are characterized by the presence of the tat basic region (amino acids 49-57), the absence of the tat cysteine-rich region (amino acids 22-36) and the absence of the tat exon 2-encoded carboxy-terminal domain (amino acids 73-86) of the naturally-occurring tat protein. By virtue of the absence of the cysteine-rich region, the preferred transport polypeptides of this invention solve the potential problems of spurious trans-activation and disulfide aggregation. The reduced size of the preferred transport polypeptides of this invention also minimizes interference with the biological activity of the cargo molecule.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L13 ANSWER 17 OF 20 USPATFULL

TITLE: • Nucleic acid conjugates of tat-derived transport polypeptides  
 INVENTOR(S): Frankel, Alan, 21 Marinero Cir. #206, Tiburon, CA, United States 94920  
 Pabo, Carl, 18 Weldon Rd., Newton, MA, United States 02158  
 Barsoum, James G., 9 Marlboro Rd., Lexington, MA, United States 02173  
 Fawell, Stephen E., One Black Horse Ter., Winchester, MA, United States 01890  
 Pepinsky, R. Blake, 30 Falmouth Rd., Arlington, MA, United States 02174

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 5670617		19970923
APPLICATION INFO.:	US 1995-450246		19950525 (8)
RELATED APPLN. INFO.:	Division of Ser. No. US 1994-235403, filed on 28 Apr 1994 which is a continuation-in-part of Ser. No. US 1993-158015, filed on 24 Nov 1993, now abandoned which is a continuation of Ser. No. US 1991-636662, filed on 2 Jan 1991, now abandoned which is a continuation-in-part of Ser. No. US 1989-454450, filed on 21 Dec 1989, now abandoned, said Ser. No. US -235403 which is a continuation-in-part of Ser. No. US 1992-934375, filed on 21 Aug 1992, now abandoned		
DOCUMENT TYPE:	Utility		
FILE SEGMENT:	Granted		
PRIMARY EXAMINER:	Elliot, George C.		
ASSISTANT EXAMINER:	Larson, Thomas G.		
LEGAL REPRESENTATIVE:	Fish & Neave, Haley, Jr., James F., Kanter, Madge R.		
NUMBER OF CLAIMS:	4		
EXEMPLARY CLAIM:	1		
NUMBER OF DRAWINGS:	21 Drawing Figure(s); 16 Drawing Page(s)		
LINE COUNT:	3849		
CAS INDEXING IS AVAILABLE FOR THIS PATENT.			

AB This invention relates to delivery of biologically active cargo molecules, such as polypeptides and nucleic acids, into the cytoplasm and nuclei of cells in vitro and in vivo. Intracellular delivery of cargo molecules according to this invention is accomplished by the use of novel transport polypeptides which comprise HIV tat protein or one or more portions thereof, and which are covalently attached to cargo molecules. The transport polypeptides in preferred embodiments of this invention are characterized by the presence of the tat basic region (amino acids 49-57), the absence of the tat cysteine-rich region (amino acids 22-36) and the absence of the tat exon 2-encoded carboxy-terminal domain (amino acids 73-86) of the naturally-occurring tat protein. By virtue of the absence of the cysteine-rich region, the preferred transport polypeptides of this invention solve the potential problems of spurious trans-activation and disulfide aggregation. The reduced size of the preferred transport polypeptides of this invention also minimizes interference with the biological activity of the cargo molecule.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L13 ANSWER 18 OF 20 USPATFULL  
 ACCESSION NUMBER: 97:66013 USPATFULL  
 TITLE: Nucleic acids encoding and methods of making tat-derived transport polypeptides  
 INVENTOR(S): Frankel, Alan, 21 Marinero Cir., Tiburon, CA, United States 94920  
 Pabo, Carl, 18 Weldon Rd., Newton, MA, United States 02158  
 Barsoum, James G., 9 Marlboro Rd., Lexington, MA, United States 02173  
 Fawell, Stephen E., One Black Horse Ter., Winchester, MA, United States 01890  
 Pepinsky, R. Blake, 30 Falmouth Rd., Arlington, MA, United States 02174

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 5652122		19970729
APPLICATION INFO.:	US 1995-450257		19950525 (8)
RELATED APPLN. INFO.:	Division of Ser. No. US 1994-235403, filed on 28 Apr 1994 which is a continuation-in-part of Ser. No. US		

is a continuation of Ser. No. US 1991-636662, filed on 2 Jan 1991, now abandoned which is a continuation-in-part of Ser. No. US 1989-454450, filed on 21 Dec 1989, now abandoned, said Ser. No. US -235403 which is a continuation-in-part of Ser. No. US 1992-934375, filed on 21 Aug 1992, now abandoned

DOCUMENT TYPE: Utility  
FILE SEGMENT: Granted  
PRIMARY EXAMINER: Elliott, George C.  
ASSISTANT EXAMINER: Larson, Thomas G.  
LEGAL REPRESENTATIVE: Fish & Neave, Haley, Jr., James F., Kanter, Madge R.  
NUMBER OF CLAIMS: 9  
EXEMPLARY CLAIM: 1  
NUMBER OF DRAWINGS: 21 Drawing Figure(s); 16 Drawing Page(s)  
LINE COUNT: 3872

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB This invention relates to delivery of biologically active cargo molecules, such as polypeptides and nucleic acids, into the cytoplasm and nuclei of cells in vitro and in vivo. Intracellular delivery of cargo molecules according to this invention is accomplished by the use of novel transport polypeptides which comprise HIV tat protein or one or more portions thereof, and which are covalently attached to cargo molecules. The transport polypeptides in preferred embodiments of this invention are characterized by the presence of the tat basic region (amino acids 49-57), the absence of the tat cysteine-rich region (amino acids 22-36) and the absence of the tat exon 2-encoded carboxy-terminal domain (amino acids 73-86) of the naturally-occurring tat protein. By virtue of the absence of the cysteine-rich region, the preferred transport polypeptides of this invention solve the potential problems of spurious trans-activation and disulfide aggregation. The reduced size of the preferred transport polypeptides of this invention also minimizes interference with the biological activity of the cargo molecule.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L13 ANSWER 19 OF 20 USPATFULL

ACCESSION NUMBER: 96:1197 USPATFULL  
TITLE: Apparatus kit for detecting Dientamoeba fragilis  
INVENTOR(S): Riordan, Neil H., 7715 E. 32nd N., Wichita, KS, United States 67226

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 5480613		19960102
APPLICATION INFO.:	US 1994-239731		19940509 (8)
RELATED APPLN. INFO.:	Division of Ser. No. US 1992-964874, filed on 22 Oct 1992, now patented, Pat. No. US 5334509, issued on 2 Aug 1994		

DOCUMENT TYPE: Utility  
FILE SEGMENT: Granted  
PRIMARY EXAMINER: Snay, Jeffrey R.  
ASSISTANT EXAMINER: wallenhorst, Maureen M.  
LEGAL REPRESENTATIVE: Hovey, Williams, Timmons & Collins  
NUMBER OF CLAIMS: 10  
EXEMPLARY CLAIM: 1  
NUMBER OF DRAWINGS: 14 Drawing Figure(s); 7 Drawing Page(s)  
LINE COUNT: 612

AB A method and apparatus for producing detectable intestinal parasites. The method includes obtaining an intestinal mucosa sample (e.g. feces) having intestinal parasites, such as Dientamoeba fragilis; and contacting the obtained intestinal mucosa sample with an acridine base compound (e.g. acridine orange and/or acridine yellow, etc.) such that the intestinal parasites become differentially stained and detectable by a human eye when viewed through a fluorescence microscope. The apparatus includes a kit or the like which includes at least one vessel or vial. Preferably, two vials are contained within the kit with one vial having an isotonic salt solution including a salt, sodium chloride, potassium phosphate, etc., and the other vial containing an acridine biological staining compound.

L13 ANSWER 20 OF 20 USPATFULL

ACCESSION NUMBER: 94:66402 USPATFULL  
TITLE: Method for detecting intestinal pathogen dientamoeba fragilis  
INVENTOR(S): Riordan, Neil H., 7715 E. 32nd North, Wichita, KS,

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 5334509		19940802
APPLICATION INFO.:	US 1992-964874		19921022 (7)
DOCUMENT TYPE:	Utility		
FILE SEGMENT:	Granted		
PRIMARY EXAMINER:	Housel, James C.		
ASSISTANT EXAMINER:	Wallenhorst, Maureen M.		
LEGAL REPRESENTATIVE:	Carpenter, John W.		
NUMBER OF CLAIMS:	20		
EXEMPLARY CLAIM:	15		
NUMBER OF DRAWINGS:	14 Drawing Figure(s); 7 Drawing Page(s)		
LINE COUNT:	681		

AB A method and apparatus for producing detectable intestinal parasites. The method includes obtaining an intestinal mucosa sample (e.g. feces) having intestinal parasites, such as *Dienamoeba fragilis*; and contacting the obtained intestinal mucosa sample with an acridine base compound (e.g. acridine orange and/or acridine yellow, etc.) such that the intestinal parasites become differentially stained and detectable by a human eye when viewed through a fluorescence microscope. The apparatus includes a kit or the like which includes at least one vessel or vial. Preferably, two vials are contained within the kit with one vial having an isotonic salt solution including a salt, such as sodium chloride, potassium phosphate, etc., and the other vial containing an acridine biological staining compound.

=> s tar and hiv and rna  
L14 3170 TAR AND HIV AND RNA

=> s l14 and nucle?  
L15 1920 L14 AND NUCLE?

=> s l14 and snorna  
L16 1 L14 AND SNORNA

=> d l16 ibib abs tot

L16 ANSWER 1 OF 1 USPATFULL  
ACCESSION NUMBER: 1999:81758 USPATFULL  
TITLE: Non-activated receptor complex proteins and uses thereof  
INVENTOR(S): Davis, Roger J., Princeton, MA, United States  
Galcheva-Gargova, Zoya, Worcester, MA, United States  
PATENT ASSIGNEE(S): University of Massachusetts, Boston, MA, United States  
(U.S. corporation)

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 5925566		19990720
APPLICATION INFO.:	US 1997-870518		19970606 (8)

	NUMBER	DATE
PRIORITY INFORMATION:	US 1996-19219P	19960606 (60)
DOCUMENT TYPE:	Utility	
FILE SEGMENT:	Granted	
PRIMARY EXAMINER:	Campell, Bruce R.	
ASSISTANT EXAMINER:	Nguyen, Dave Trong	
LEGAL REPRESENTATIVE:	Fish & Richardson, P.C.	
NUMBER OF CLAIMS:	23	
EXEMPLARY CLAIM:	1	
NUMBER OF DRAWINGS:	22 Drawing Figure(s); 18 Drawing Page(s)	
LINE COUNT:	2438	

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB The invention features a substantially pure ZPR1 polypeptide. For example, a ZPR1 polypeptide that specifically binds to a non-activated membrane-bound receptor (e.g., EGF or PDGF receptors) and specifically binds small nucleolar RNAs (e.g., U3). ZPR1 polypeptides can be isolated from any eukaryote, including mammals (e.g. rodents and humans) and fungi (e.g., *S. cerevisiae* and *S. pombe*).

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

=> s 114 and (small nucleolar)  
L17 7 L14 AND (SMALL NUCLEOLAR)

=> s 117 not 116  
L18 6 L17 NOT L16

=> d 118 ibib abs tot

L18 ANSWER 1 OF 6 MEDLINE  
ACCESSION NUMBER: 2002647867 MEDLINE  
DOCUMENT NUMBER: 22294961 PubMed ID: 12376617  
TITLE: A nucleolar \*\*\*TAR\*\*\* decoy inhibitor of \*\*\*HIV\*\*\*  
-1 replication.  
AUTHOR: Michienzi Alessandro; Li Shirley; Zaia John A; Rossi John J  
CORPORATE SOURCE: Divisions of Molecular Biology and Virology, Beckman  
Research Institute of the City of Hope, 1450 East Duarte  
Road, Duarte, CA 91010-3011, USA.  
CONTRACT NUMBER: AI29329 (NIAID)  
AI46030 (NIAID)  
SOURCE: PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE  
UNITED STATES OF AMERICA, (2002 Oct 29) 99 (22) 14047-52.  
Journal code: 7505876. ISSN: 0027-8424.  
PUB. COUNTRY: United States  
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 200212  
ENTRY DATE: Entered STN: 20021031  
Last Updated on STN: 20021217  
Entered Medline: 20021209

AB Tat is a critical regulatory factor in \*\*\*HIV\*\*\* -1 gene expression. It  
mediates the transactivation of transcription from the \*\*\*HIV\*\*\* -1 LTR  
by binding to the transactivation response ( \*\*\*TAR\*\*\* ) element in a  
complex with cyclin T1. Because of its critical and early role in  
\*\*\*HIV\*\*\* gene expression, Tat and its interaction with the \*\*\*TAR\*\*\*  
element constitute important therapeutic targets for the treatment of  
\*\*\*HIV\*\*\* -1 infection. Based on the known nucleolar localization  
properties of Tat, we constructed a chimeric \*\*\*small\*\*\*  
\*\*\*nucleolar\*\*\* \*\*\*RNA\*\*\* - \*\*\*TAR\*\*\* decoy that localizes to the  
nucleoli of human cells and colocalizes in the nucleolus with a  
Tat-enhanced GFP fusion protein. When the chimeric \*\*\*RNA\*\*\* was  
stably expressed in human T lymphoblastoid CEM cells it potently inhibited  
\*\*\*HIV\*\*\* -1 replication. These results demonstrate that the nucleolar  
trafficking of Tat is critical for \*\*\*HIV\*\*\* -1 replication and  
suggests a role for the nucleolus in \*\*\*HIV\*\*\* -1 viral replication.

L18 ANSWER 2 OF 6 CAPLUS COPYRIGHT 2002 ACS  
ACCESSION NUMBER: 2002:859140 CAPLUS  
DOCUMENT NUMBER: 137:336616  
TITLE: A nucleolar \*\*\*TAR\*\*\* decoy inhibitor of  
\*\*\*HIV\*\*\* -1 replication  
AUTHOR(S): Michienzi, Alessandro; Li, Shirley; Zaia, John A.;  
Rossi, John J.  
CORPORATE SOURCE: Division of Molecular Biology, Beckman Research  
Institute of the City of Hope, Duarte, CA, 91010-3011,  
USA  
SOURCE: Proceedings of the National Academy of Sciences of the  
United States of America (2002), 99(22), 14047-14052  
CODEN: PNASA6; ISSN: 0027-8424  
PUBLISHER: National Academy of Sciences  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB Tat is a crit. regulatory factor in \*\*\*HIV\*\*\* -1 gene expression. It  
mediates the transactivation of transcription from the \*\*\*HIV\*\*\* -1 LTR  
by binding to the transactivation response ( \*\*\*TAR\*\*\* ) element in a  
complex with cyclin T1. Because of its crit. and early role in  
\*\*\*HIV\*\*\* gene expression, Tat and its interaction with the \*\*\*TAR\*\*\*  
element constitute important therapeutic targets for the treatment of  
\*\*\*HIV\*\*\* -1 infection. Based on the known nucleolar localization  
properties of Tat, the authors constructed a chimeric \*\*\*small\*\*\*  
\*\*\*nucleolar\*\*\* \*\*\*RNA\*\*\* - \*\*\*TAR\*\*\* decoy that localizes to the  
nucleoli of human cells and colocalizes in the nucleolus with a  
Tat-enhanced GFP fusion protein. When the chimeric \*\*\*RNA\*\*\* was  
stably expressed in human T lymphoblastoid CEM cells it potently inhibited  
\*\*\*HIV\*\*\* -1 replication. These results demonstrate that the nucleolar  
trafficking of Tat is crit. for \*\*\*HIV\*\*\* -1 replication and suggests a

REFERENCE COUNT: 52 THERE ARE 52 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L18 ANSWER 3 OF 6 LIFESCI COPYRIGHT 2002 CSA  
ACCESSION NUMBER: 2002:119753 LIFESCI  
TITLE: A nucleolar \*\*\*TAR\*\*\* decoy inhibitor of \*\*\*HIV\*\*\*  
-1 replication  
AUTHOR: Michienzi, A.; Li, S.; Zaia, J.A.; Rossi, J.J.  
CORPORATE SOURCE: Divisions of Molecular Biology and Virology, Beckman  
Research Institute of the City of Hope, 1450 East Duarte  
Road, Duarte, CA 91010-3011; E-mail: jrossi@bricoh.edu  
SOURCE: Proceedings of the National Academy of Sciences, USA [Proc.  
Nat'l. Acad. Sci. USA], (20021029) vol. 99, no. 22, pp.  
14047-14052.  
ISSN: 0027-8424.  
DOCUMENT TYPE: Journal  
FILE SEGMENT: N; V  
LANGUAGE: English  
SUMMARY LANGUAGE: English  
AB Tat is a critical regulatory factor in \*\*\*HIV\*\*\* -1 gene expression. It  
mediates the transactivation of transcription from the \*\*\*HIV\*\*\* -1 LTR  
by binding to the transactivation response ( \*\*\*TAR\*\*\* ) element in a  
complex with cyclin T1. Because of its critical and early role in  
\*\*\*HIV\*\*\* gene expression, Tat and its interaction with the \*\*\*TAR\*\*\*  
element constitute important therapeutic targets for the treatment of  
\*\*\*HIV\*\*\* -1 infection. Based on the known nucleolar localization  
properties of Tat, we constructed a chimeric \*\*\*small\*\*\*  
\*\*\*nucleolar\*\*\* \*\*\*RNA\*\*\* - \*\*\*TAR\*\*\* decoy that localizes to the  
nucleoli of human cells and colocalizes in the nucleolus with a  
Tat-enhanced GFP fusion protein. When the chimeric \*\*\*RNA\*\*\* was  
stably expressed in human T lymphoblastoid CEM cells it potently inhibited  
\*\*\*HIV\*\*\* -1 replication. These results demonstrate that the nucleolar  
trafficking of Tat is critical for \*\*\*HIV\*\*\* -1 replication and  
suggests a role for the nucleolus in \*\*\*HIV\*\*\* -1 viral replication.

L18 ANSWER 4 OF 6 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.  
ACCESSION NUMBER: 2002394185 EMBASE  
TITLE: A nucleolar \*\*\*TAR\*\*\* decoy inhibitor of \*\*\*HIV\*\*\*  
-1 replication.  
AUTHOR: Michienzi A.; Li S.; Zaia J.A.; Rossi J.J.  
CORPORATE SOURCE: J.J. Rossi, Divisions of Molecular Biology, Beckman Res.  
Inst. the City of Hope, 1450 East Duarte Road, Duarte, CA  
91010-3011, United States. jrossi@bricoh.edu  
SOURCE: Proceedings of the National Academy of Sciences of the  
United States of America, (29 Oct 2002) 99/22  
(14047-14052).  
Refs: 52  
ISSN: 0027-8424 CODEN: PNASA6  
COUNTRY: United States  
DOCUMENT TYPE: Journal; Article  
FILE SEGMENT: 004 Microbiology  
030 Pharmacology  
037 Drug Literature Index  
LANGUAGE: English  
SUMMARY LANGUAGE: English  
AB Tat is a critical regulatory factor in \*\*\*HIV\*\*\* -1 gene expression. It  
mediates the transactivation of transcription from the \*\*\*HIV\*\*\* -1 LTR  
by binding to the transactivation response ( \*\*\*TAR\*\*\* ) element in a  
complex with cyclin T1. Because of its critical and early role in  
\*\*\*HIV\*\*\* gene expression, Tat and its interaction with the \*\*\*TAR\*\*\*  
element constitute important therapeutic targets for the treatment of  
\*\*\*HIV\*\*\* -1 infection. Based on the known nucleolar localization  
properties of Tat, we constructed a chimeric \*\*\*small\*\*\*  
\*\*\*nucleolar\*\*\* \*\*\*RNA\*\*\* - \*\*\*TAR\*\*\* decoy that localizes to the  
nucleoli of human cells and colocalizes in the nucleolus with a  
Tat-enhanced GFP fusion protein. When the chimeric \*\*\*RNA\*\*\* was  
stably expressed in human T lymphoblastoid CEM cells it potently inhibited  
\*\*\*HIV\*\*\* -1 replication. These results demonstrate that the nucleolar  
trafficking of Tat is critical for \*\*\*HIV\*\*\* -1 replication and  
suggests a role for the nucleolus in \*\*\*HIV\*\*\* -1 viral replication.

L18 ANSWER 5 OF 6 USPATFULL  
ACCESSION NUMBER: 2002:217408 USPATFULL  
TITLE: Oligomers that bind to ku protein  
INVENTOR(S): Dynan, William S., Augusta, GA, United States  
Yoo, Sunghan, Augusta, GA, United States

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 6441158	B1	20020827
APPLICATION INFO.:	US 1998-223139		19981230 (9)

	NUMBER	DATE
PRIORITY INFORMATION:	US 1997-70278P	19971231 (60)
DOCUMENT TYPE:	Utility	
FILE SEGMENT:	GRANTED	
PRIMARY EXAMINER:	Wang, Andrew	
LEGAL REPRESENTATIVE:	Kilpatrick Stockton LLP, Rothschild, Cynthia B., Calkins, Charles W.	
NUMBER OF CLAIMS:	8	
EXEMPLARY CLAIM:	1	
NUMBER OF DRAWINGS:	8 Drawing Figure(s); 5 Drawing Page(s)	
LINE COUNT:	2333	

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB Disclosed are oligomers that bind Ku protein. These oligomers are useful for inhibiting activation of DNA-PK, treating certain forms of autoimmune disease, detection and purification of Ku protein, and identification of proteins that interact with Ku protein. Preferably, the oligomers are composed of nucleotides, nucleotide analogs, or a combination. Most preferably, the oligomers are composed of ribonucleotides. Also disclosed is a method of inhibiting DNA repair, a method of identifying cellular proteins that interact with Ku protein, and a method of treating autoimmune disease in patients with anti-Ku antibodies. The disclosed oligomers can have several preferred features, either alone or in combination, in addition to Ku binding. One such feature, referred to herein as inhibition activity, is inhibition of DNA-PK kinase activity. Another preferred feature, referred to herein as aptamer motifs, is the presence of one or more of the base sequences GCUUUGCCANNAC, A(A/C)AUGA, and AACUUGCA. These sequences--referred to herein as aptamer motif 1, aptamer motif 2, and aptamer motif 3, respectively--are associated with Ku binding capability. Another preferred feature, referred to herein as aptamer structure, is the presence of a structure similar to the structure shown in FIG. 6A. This structure has the general formula 5'-A-B-C-D-C'-E-A'-3', where A, B, C, D, C', E, and A' are components of the oligomer. In this structure, A and A' interact to form a stem structure, C and C' interact to form a stem structure, B and E make up a bulge region, and D is either a bulge or a loop. FIG. 6A depicts component D as a loop. Each of these preferred features (inhibition activity, aptamer motif, and aptamer structure) can be used either alone or in combination with one or both of the other characteristics.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L18 ANSWER 6 OF 6 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

ACCESSION NUMBER: 2003:15084 BIOSIS

DOCUMENT NUMBER: PREV200300015084

TITLE: A nucleolar \*\*\*TAR\*\*\* decoy inhibitor of \*\*\*HIV\*\*\*  
-1 replication.AUTHOR(S): Michienzi, Alessandro; Li, Shirley; Zaia, John A.; Rossi,  
John J. (1)CORPORATE SOURCE: (1) Divisions of Molecular Biology, Beckman Research  
Institute of the City of Hope, 1450 East Duarte Road,  
Duarte, CA, 91010-3011, USA: jrossi@bricoh.edu USASOURCE: Proceedings of the National Academy of Sciences of the  
United States of America, (October 29 2002) vol. 99, No.  
22, pp. 14047-14052. print.  
ISSN: 0027-8424.

DOCUMENT TYPE: Article

LANGUAGE: English

AB Tat is a critical regulatory factor in \*\*\*HIV\*\*\* -1 gene expression. It mediates the transactivation of transcription from the \*\*\*HIV\*\*\* -1 LTR by binding to the transactivation response ( \*\*\*TAR\*\*\* ) element in a complex with cyclin T1. Because of its critical and early role in \*\*\*HIV\*\*\* gene expression, Tat and its interaction with the \*\*\*TAR\*\*\* element constitute important therapeutic targets for the treatment of \*\*\*HIV\*\*\* -1 infection. Based on the known nucleolar localization properties of Tat, we constructed a chimeric \*\*\*small\*\*\*  
\*\*\*nucleolar\*\*\* \*\*\*RNA\*\*\* - \*\*\*TAR\*\*\* decoy that localizes to the nucleoli of human cells and colocalizes in the nucleolus with a



- stably expressed in human T lymphoblastoid CEM cells it potently inhibited  
\*\*\*HIV\*\*\* -1 replication. These results demonstrate that the nucleolar  
trafficking of Tat is critical for \*\*\*HIV\*\*\* -1 replication and  
suggests a role for the nucleolus in \*\*\*HIV\*\*\* -1 viral replication.

=> d 5 kwic

L18 ANSWER 5 OF 6 USPATFULL

SUMM There is some evidence that Ku protein interacts with \*\*\*RNA\*\*\* ,  
although this has been much less studied than the interaction with DNA.  
Antibodies to Ku protein stain both the nucleoplasm. . . not (Reeves,  
J. Exp. Med. 161, 18-39 (1985)). Thus, nucleolar localization may be  
regulated by interaction of Ku protein with \*\*\*RNA\*\*\* . Ku protein  
does not appear to bind to bulk tRNA or to synthetic \*\*\*RNA\*\*\*  
polymers (Mimori & Hardin, J. Biol. Chem. 261, 10375-10379 (1986)).  
However, one study showed that Ku protein forms a specific complex with  
an \*\*\*RNA\*\*\* that included the \*\*\*HIV\*\*\* trans-activation  
response ( \*\*\*TAR\*\*\* ) element sequence (Kaczmariski & Khan, Biochem.  
Biophys. Res. Commun. 196, 935-942 (1993)).

DRWD FIG. 1 is a graph of percent \*\*\*RNA\*\*\* or DNA bound versus  
concentration of Ku (nM) for five representative Ku protein- \*\*\*RNA\*\*\*  
and DNA binding curves obtained with an electrophoretic mobility shift  
assay. The graph shows that pooled \*\*\*RNA\*\*\* tested after the fourth  
and sixth round of Systematic Evolution of Ligands by Exponential  
Enrichment (SELEX; Tuerk & Gold, Science. . . and 5,567,588) had an  
increased ability to bind to Ku protein. Radiolabeled nucleic acid  
probes were as follows: .quadrature., nonselected \*\*\*RNA\*\*\* ;  
.gradient., \*\*\*RNA\*\*\* after 4th round of selection;  
.tangle-solidup., \*\*\*RNA\*\*\* after 6th round of selection; 0,  
\*\*\*HIV\*\*\* - \*\*\*TAR\*\*\* \*\*\*RNA\*\*\* (included for comparison);  
.diamond-solid., 21 base pair double-stranded DNA.

DRWD FIG. 2A is a graph of kinase activity (percentage of activity in absence  
of \*\*\*RNA\*\*\* ) versus \*\*\*RNA\*\*\* concentration (nM) for eight  
representative aptamers. The aptamers were as follows: X, nonselected  
\*\*\*RNA\*\*\* ; 0, #2; .quadrature., #7-3; .DELTA., SC4; +, SC5;  
.diamond-solid., SC8; .tangle-soliddn., #1-2; .circle-solid., SC2.

DRWD FIG. 2B is a graph of DNA binding (percent of Ku-DNA complexes detected  
in the absence of competitor) versus \*\*\*RNA\*\*\* concentration (nM)  
for seven representative aptamers. Purified Ku protein (2 nM) was  
incubated with radiolabeled 21 base pair double-stranded DNA (1 nM) in  
the presence of various amounts of nonselected \*\*\*RNA\*\*\* or aptamer.  
The aptamers were as follows: X, nonselected \*\*\*RNA\*\*\* ; 0, #2;  
.quadrature., #7-3; .DELTA., SC4; +, SC5; .diamond-solid., SC8;  
.tangle-soliddn., #1-2; .circle-solid., SC2.

DRWD FIG. 3 is a bar graph of kinase activity (percentage of activity in the  
absence of \*\*\*RNA\*\*\* ) in the presence of different aptamers, each at  
three different concentrations. Reactions were performed in crude  
nuclear extract. Reactions contained. . .

DRWD . . . '3, SEQ ID NOS: 22, 6, and 23) and #2 ('5 to '3, SEQ ID NOS:  
22, 5, and 23). \*\*\*RNA\*\*\* secondary structures were predicted using  
the methods of Zuker and co-workers (Jaeger et al., 1989; Jaeger et al.,  
1990) as. . .

DETD . . . described in the examples. Only the portion of the selected  
oligomers that arose from the variable region of the pool \*\*\*RNA\*\*\*  
is shown in Table 1. That is, each of the oligonucleotides listed in  
Table 1 include the sequence GGGAGGAUUAUUUCAGACCGUAA (SEQ ID. . .

DETD  
TABLE 1

\*\*\*RNA\*\*\* Sequences and Kd values from the SELEX Procedure

Name DNA-PK  
Frequency Aligned sequence Inhibition K.sub.d

CLASS I

SC6 (3) GACUCACGAUGGACCAUACGCCUCCCCACUGGUCUUGUUA (SEQ ID. . . ID NO:17) A  
0.8 nM

#42 (1) CAUCCUGGUACUCACUUCGACAUCGUACGUUCAAUCAUAC (SEQ. B) NO:18) B 4.5 nM

OTHER

SC13 (3) ACCUUUUUAGACGAACCUCAAAGUACAUUUAGUUGAAAAC (SEQ ID NO:19) B 0.8 nM

\*\*\*RNA\*\*\* sequences are aligned to show maximum sequence identity. The  
underlined sequence indicates sequence identity within class.

DETD \*\*\*RNA\*\*\* is relatively labile and can be degraded by a number of  
ribonucleases. This degradation can be greatly reduced by the. . .  
and substitutions at the 2'-prime position of the ribonucleotide and by

\*\*\*RNA\*\*\* . In addition a variety of modifications can be made on the nucleobases themselves which both inhibit degradation and which can.

DETD In preferred oligomers, the oligonucleotides are made from \*\*\*RNA\*\*\* . The examples of Ku aptamers obtained as described in the examples are in the form of \*\*\*RNA\*\*\* . These RNAs were selected from a pool of about 10<sup>sup</sup>.14 different \*\*\*RNA\*\*\* sequences using the systematic evolution of ligands by exponential enrichment (SELEX) procedure. Most of the selected RNAs bind to Ku.

DETD . . . the behavior of molecules with each other. These applications can be adapted to define and display the secondary structure of \*\*\*RNA\*\*\* and DNA molecules.

DETD . . . be synthesized by solid phase .beta.-cyanoethyl phosphoramidite chemistry (Sinha et al., Nucleic Acids Res. 12:4539-4557 (1984)) on any commercially available DNA/ \*\*\*RNA\*\*\* synthesizer. A preferred method is the 2'-O-tert-butyl dimethylsilyl (TBDMS) protection strategy for the ribonucleotides (Usman et al., J. Am. Chem. Soc. . . . its advantageous properties (McCollum and Andrus Tetrahedron Letters 32:4069-4072 (1991)). Fluorescein can be added to the 5'-end of a substrate \*\*\*RNA\*\*\* during the synthesis by using commercially available fluorescein phosphoramidites. In general, a desired oligomer can be synthesized using a standard \*\*\*RNA\*\*\* cycle. Upon completion of the assembly, all base labile protecting groups are removed by an 8 hour treatment at 55.degree..

DETD . . . oligomers can also be produced through enzymatic methods, when the nucleotide subunits are available for enzymatic manipulation. For example, the \*\*\*RNA\*\*\* molecules can be made through in vitro \*\*\*RNA\*\*\* polymerase T7 reactions. They can also be made by strains of bacteria or cell lines expressing T7, and then subsequently.

DETD . . . viral vectors are Adenovirus, Adeno-associated virus, Herpes virus, Vaccinia virus, Polio virus, AIDS virus, neuronal trophic virus, Sindbis and other \*\*\*RNA\*\*\* viruses, including these viruses with the \*\*\*HIV\*\*\* backbone. Also preferred are any viral families which share the properties of these viruses which make them suitable for use.

DETD . . . chemical or physical methods to introduce genes into cells. Typically, viral vectors contain, nonstructural early genes, structural late genes, an \*\*\*RNA\*\*\* polymerase III transcript, inverted terminal repeats necessary for replication and encapsidation, and promoters to control the transcription and replication of.

DETD . . . a primer binding site to bind the tRNA primer of reverse transcription, terminal repeat sequences that guide the switch of \*\*\*RNA\*\*\* strands during DNA synthesis, a purine rich sequence 5' to the 3' LTR that serve as the priming site for.

DETD . . . relatively fixed location in regard to the transcription start site. A promoter contains core elements required for basic interaction of \*\*\*RNA\*\*\* polymerase and transcription factors, and may contain upstream elements and response elements.

DETD Expression of the disclosed aptamers in cells also requires sequences to direct expression. It is preferred that an \*\*\*RNA\*\*\* polymerase III (pol III) promoter be used for expression. Pol III promoters generate transcripts that can be engineered to remain. . . . capping signal, and termination sequence. Pol III promoters, and other pol III transcription signals, are present in tRNA genes, 5S \*\*\*RNA\*\*\* genes, small nuclear \*\*\*RNA\*\*\* genes, and small cytoplasmic \*\*\*RNA\*\*\* genes. Preferred pol III promoters for use in aptamer expression vectors are the human small nuclear U6 gene promoter and tRNA gene promoters. The use of U6 gene transcription signals to produce short \*\*\*RNA\*\*\* molecules in vivo is described by Noonberg et al., Nucleic Acids Res. 22:2830-2836 (1994), and the use of tRNA transcription.

DETD . . . Biol. Chem. 268:7868-7873 (1993), and Romero and Blackburn, Cell 67:343-353 (1991). The use of pol III promoters for expression of \*\*\*RNA\*\*\* molecules is also described in WO 95/23225 by Ribozyme Pharmaceuticals, Inc.

DETD \*\*\*RNA\*\*\* fusion constructs can be produced which aid in the correct and specific targeting of expressed \*\*\*RNA\*\*\* molecules. These types of \*\*\*RNA\*\*\* fusion constructs are discussed in detail in Good et al., "Expression of Small Therapeutic RNAs in Cell Nuclei", Gene Therapy, 4:45-54 (1997). Expression vectors based on human tRNA(met) and U6 snRNA promoters are useful for targeting \*\*\*RNA\*\*\* expression and to render the resulting \*\*\*RNA\*\*\* transcripts more resistant to degradation.

DETD . . . coupled to the oligomer. As used herein, a capture tag is any compound that can be associated with a synthesized \*\*\*RNA\*\*\* molecule and which can be used to separate compounds or complexes having the capture tag from those that do not.

coupled to the 5' end of the synthesized \*\*\*RNA\*\*\* molecule. Preferred capture tags, described in the context of nucleic acid probes, have been described by Syvanen et al., Nucleic.

DETD . . . or bind other groups of antibodies. Thus, one can have an antibody to the capture tag and then this antibody:capture tag: \*\*\*RNA\*\*\* complex can then be purified by binding to an antibody for the antibody portion of the complex.

DETD . . . molecules of choice. For example, a preferred capture tag of this type is one which contains a sulfur atom. An \*\*\*RNA\*\*\* molecule which is associated with this capture tag can be purified by retention on a thiolpropyl sepharose column. Extensive washing of the column removes unwanted molecules and reduction with .beta.-mercaptoethanol, for example, allows the desired \*\*\*RNA\*\*\* molecules to be collected after purification under relatively gentle conditions (See Lorsch and Szostak, 1994 for a reduction to practice).

DETD B. Identification Of Cellular \*\*\*RNA\*\*\* Sequences Predicted That Bind Ku Protein

DETD . . . bind tightly to Ku protein is consistent with suggestions that Ku protein function may be regulated to some extent by \*\*\*RNA\*\*\* in the normal cell. (Reeves, J. Exp. Med. 161, 18-39 (1985)). One site where such interactions might occur is in. . . Med. 161, 18-39 (1985)). A number of studies have shown that DNA-PK and Ku protein have an ability to regulate \*\*\*RNA\*\*\* polymerase I, which is localized in the nucleolus (Hoff et al., Proc. Natl. Acad. Sci. USA 91, 762-766 (1994); Kuhn. . . however, Ku protein associates with nucleolar RNAs, this would provide a potential targeting mechanism. Nucleoli contain a multitude of discrete \*\*\*small\*\*\* \*\*\*nucleolar\*\*\* RNAs, as well as nascent ribosomal \*\*\*RNA\*\*\* (reviewed in Smith & Steitz, Cell 89, 669-672 (1997)). The consensus sequences present in the aptamers can be used to predict whether particular nucleolar RNAs or other cellular RNAs are likely to bind to Ku protein. Cellular \*\*\*RNA\*\*\* sequences that become available in public databases can be searched to determine whether there is identity with the consensus sequences.

DETD . . . acid-like properties, such as sequence dependent hybridization, that contain at one or more positions, a modification away from a standard \*\*\*RNA\*\*\* or DNA nucleotide. A preferred example of an oligonucleotide analogue is peptide nucleic acid.

DETD Preparing the \*\*\*RNA\*\*\* Pool and Screening for Ku Protein Binding

DETD Because there was little previous information about Ku protein-\*\*\*RNA\*\*\* interactions, the \*\*\*RNA\*\*\* binding properties of Ku protein were systematically investigated using SELEX (systematic evolution of ligands by exponential enrichment) technology. with this.

DETD The \*\*\*RNA\*\*\* pool was prepared by synthesis of single stranded \*\*\*RNA\*\*\* from a template comprising a region of conserved sequences and a region of randomized and/or biased sequences.

DETD Synthesis of the \*\*\*RNA\*\*\* Pool

DETD SELEX protocols and the template used to synthesize the nonselected \*\*\*RNA\*\*\* pool were obtained from Dr. Hang Chen and Dr. Larry Gold (University of Colorado) and modified as described below. The. . .

DETD . . . carried out largely as described in Chen & Gold, Biochemistry 33, 8746-8756 (1994). The template for synthesis of the starting \*\*\*RNA\*\*\* pool was based on a DNA oligonucleotide, 5'-CCCGGATCCTAGTTCACGATGCTGCAA-(N).sub.40-TTACGGTCTGAGAAAATATCCTCCC-3'(SEQ ID NO:27), where N indicates an equimolar mixture of A, G, C. .

DETD This template preparation was incubated for 2-3 h at 37.degree. C. in a reaction mixture containing 475 U/ml T7 \*\*\*RNA\*\*\* polymerase, 40 mM Tris-HCl, pH 8.0, 12 mM MgCl.sub.2, 5 mM DTT, 1 mM spermidine, 0.002% Triton X-100, 4% PEG. . . was continued for 30 minute. The resulting RNAs were fractionated by 10% urea-PAGE and gel slices containing full-length 92 nucleotide \*\*\*RNA\*\*\* were excised, crushed, mixed with 0.5 ml TE, 0.5 ml phenol, and 10 .mu.l 10% sodium dodecyl sulfate, and tumbled overnight at 4.degree. C. to extract \*\*\*RNA\*\*\*. The supernatant was collected and \*\*\*RNA\*\*\* was precipitated with NaOAc and EtOH as described above. \*\*\*RNA\*\*\* was dissolved in TE, denatured at 100.degree. C. for 2 minute, and renatured by adjusting to 5 mM MgCl.sub.2, cooling rapidly to 0.degree. C., and incubating for 30 minute. The amount of \*\*\*RNA\*\*\* recovered was estimated by liquid scintillation counting.

DETD To perform in vitro selections, the \*\*\*RNA\*\*\* pool was mixed with purified Ku protein under conditions for complex binding. \*\*\*RNA\*\*\* and Ku protein were incubated for 30 minute at room temperature in a buffer containing 25 mM Tris-HCl, pH 7.9, . . . glycerol, 5 mM MgCl.sub.2, 0.5 mM DTT, and 0.01% Tween 20 in a final volume of 50-100 .mu.l. Concentrations of \*\*\*RNA\*\*\*, Ku protein, and KCl were

were isolated by electrophoresis on a 5% non-denaturing polyacrylamide gel containing 25 mM Tris-HCl, pH 8.3, 190 mM glycine, . . . over a nitrocellulose filter (Millipore 25- $\mu$ m HAWP), which was washed twice with 5 ml of 25 mM Tris-HCl, pH 7.9. \*\*\*RNA\*\*\* was eluted from the polyacrylamide gel or the filter by tumbling in a mixture of TE, phenol, and SDS as.

DETD Once the enriched \*\*\*RNA\*\*\* molecules were separated from the isolated complexes, they were amplified by reverse transcription followed by polymerase chain reaction. The complexity.

DETD When additional rounds of selection were performed, purified Ku protein (from HeLa cell nuclear extracts) was mixed with the \*\*\*RNA\*\*\* pool and allowed to bind. Ku- \*\*\*RNA\*\*\* complexes were isolated using either a nitrocellulose filter binding or an electrophoretic mobility shift assay (EMSA). \*\*\*RNA\*\*\* that was bound to Ku protein was isolated, amplified by RT-PCR, and again enriched based on the ability to bind to Ku protein. The selected \*\*\*RNA\*\*\* was reverse-transcribed by incubating in a reaction mixture containing 50 mM Tris-HCl, pH 8.3, 60 mM NaCl, 6 mM Mg(OAc)<sub>2</sub>, . . . Selection conditions used for each round of selection are given in Table 2. To maintain the stringency of selection, the \*\*\*RNA\*\*\*:protein ratio, the KCl concentration, or both were progressively increased as shown in Table 2. Two independent selections were conducted using different batches of starting \*\*\*RNA\*\*\*.

DETD After six or seven rounds of selection, cDNA was synthesized from the final \*\*\*RNA\*\*\* pool and amplified by 5 cycles of PCR using primer 1 and 2. The product was digested with Bam HI.

DETD

## TABLE 2

### Summary of the Selection Parameters

#### Selection I

SELEX Input	***RNA***	Ku protein	sup.a	KCl	sup.b	Selection	sup.c
Round (nM)	(nM)	(mM)	Method				
1	1305	64.1	50	NCFA			
2	612.5	24.3	50	EMSA			
				EMSA			
4	1130	42	120	EMSA			
5	2320	64	150	EMSA			
6	142	3.2	160	EMSA			

#### Selection II

SELEX Input	***RNA***	Ku protein	KCl	Selection
Round (nM)	(nM)	(mM)	Method	
1	1330	53	120	EMSA
2	1660	42	120	EMSA
				160 EMSA
6	2710	18.6	160	EMSA
7	436	2.4	160	EMSA

.sup.aThe ratio of Ku protein to input \*\*\*RNA\*\*\* was gradually decreased in order to increase the stringency.

.sup.bSalt concentration was increased to maintain high stringency.

.sup.cNitrocellulose.

DETD . . . PK enzymatic activity. The ability of Ku protein to bind tightly to specific RNAs is consistent with a role for \*\*\*RNA\*\*\* in the regulation of Ku protein activity or intranuclear localization. Additionally, the identification of diverse RNAs that bind avidly to.

DETD Binding of \*\*\*RNA\*\*\* and DNA to Ku protein was measured using EMSA as shown by the representative Ku protein- \*\*\*RNA\*\*\* binding curves after successive rounds of SELEX in FIG. 1. Binding curves were obtained with EMSA followed by a nonlinear least squares fit of the data. The \*\*\*RNA\*\*\* pools in FIG. 1 were transcribed from templates prior to the first round (nonselected), and after the fourth (4th), and the sixth (6th) rounds of SELEX using T7 \*\*\*RNA\*\*\* polymerase as described in example 2. A twenty-one base pair double-stranded DNA was end-labeled with T4 polynucleotide kinase. The DNA.

DETD Ku protein bound to nonselected \*\*\*RNA\*\*\* with an apparent average K<sub>sub.d</sub> of 24 nM. This binding was somewhat stronger than expected, given that previous work had found that Ku protein had little ability to interact with tRNA or synthetic \*\*\*RNA\*\*\* polymers (Mimori & Hardin, J. Biol. Chem. 261, 10375-10379 (1986)). Pooled \*\*\*RNA\*\*\* tested after the 4th and 6th round of SELEX showed an increased ability to bind

bound with an apparent  $K_{sub.d}$  of about 0.3 nM, which was comparable to the binding seen with a double-stranded DNA oligonucleotide under similar conditions. Although the average affinity of the selected \*\*\*RNA\*\*\* was only 75-fold greater than for the nonselected pool, subsequent experiments revealed clear functional differences, as only the selected RNAs.

DETD Individual aptamer RNAs were synthesized by T7 \*\*\*RNA\*\*\* polymerase using linearized plasmid template. Both radiolabeled \*\*\*RNA\*\*\* and nonradiolabeled \*\*\*RNA\*\*\* were prepared using a MEGA shortscript T7 kit (Ambion) with 8  $\mu$ g linearized plasmid template. The \*\*\*RNA\*\*\* was gel-purified, heated, and refolded as described in example 1, and the final concentration was determined spectrophotometrically. Secondary structures were. . . Acad. Sci. USA 86, 7706-7710 (1989); Jaeger et al., Meth. Enzymol. 183, 281-306 (1990)) as implemented on the mfold server [www.ibc.wustl.edu/%7Ezucker/](http://www.ibc.wustl.edu/%7Ezucker/) \*\*\*rna\*\*\* /form 1.cgi).

DETD Synthesis of \*\*\*RNA\*\*\* containing the \*\*\*TAR\*\*\* Sequence

DETD Pools of \*\*\*RNA\*\*\* at various stages of selection were also characterized directly for their ability to bind to Ku protein in an electrophoretic. . . mobility shift assay. For comparison of the aptamer RNAs binding affinity with other known RNAs, these assays also included a \*\*\*HIV\*\*\* \*\*\*TAR\*\*\* \*\*\*RNA\*\*\*, which was synthesized by T7 \*\*\*RNA\*\*\* polymerase as described in the preceding section. This \*\*\*RNA\*\*\* consisted of the following \*\*\*HIV\*\*\*-derived sequence: GGGUCUCUCUGGUUAGACCAGAUCCUGGAGCCUGGAGCUCUCUGGC UAACUAGGGAACCC (SEQ ID NO:31).

DETD \*\*\*TAR\*\*\*-containing RNAs have previously been reported to bind selectively to Ku protein (Kaczmarek & Khan, Biochem. Biophys. Res. Commun. 196, 935-942. . .

DETD Binding of the \*\*\*TAR\*\*\* \*\*\*RNA\*\*\* was measurable, but weak, under the conditions of the experiments. These results confirm that \*\*\*TAR\*\*\*-containing \*\*\*RNA\*\*\* has some ability to bind to Ku protein, but indicate that the binding is not particularly strong, relative to other. . .

DETD After six and seven rounds of SELEX (Selection I and II in Table 2, respectively), the pooled \*\*\*RNA\*\*\* was reverse-transcribed, PCR-amplified, and cloned into plasmid vectors. 82 clones were isolated, including 63 from Selection I and 19 from. . .

DETD . . . designated SEQ ID NO:20) (wherein N may be A, C, G or T), which was perfectly conserved in three independent \*\*\*RNA\*\*\* sequences and partially conserved in four others. A second motif, AMAUGA (herein designated SEQ ID NO:21) (wherein M may be. . . partially overlaps the second, had the sequence AACUUCGA (bases 31-38 of SEQ ID NO:15). This motif was present in one \*\*\*RNA\*\*\* and partially conserved in four others. Several RNAs fell into hybrid classes containing two of the three motifs. Subsequent analysis. . .

DETD Inhibition of DNA-PK Activity by Different \*\*\*RNA\*\*\* Aptamers

DETD Binding of the \*\*\*RNA\*\*\* aptamers to Ku protein to determine if they are capable of regulating Ku protein activity was then measured. One of. . .

DETD . . . block the ability of Ku protein to activate DNA-PK. Each of the nineteen selected aptamers, as well as the nonselected \*\*\*RNA\*\*\* pool, was tested in a DNA-PK assay. Each \*\*\*RNA\*\*\* was prepared by in vitro transcription and added to a DNA-PK phosphorylation assay.

DETD . . . (EPPLSQEAFADLLWKK (SEQ ID NO:32)) containing a DNA-PK phosphorylation site from p53 (underlined), and the indicated amounts (8, 16, 32 nM) of \*\*\*RNA\*\*\* aptamer. Native DNA-PKs were purified from HeLa cell nuclear extracts as described in example 2. In FIG. 2A, DNA-PK activity is expressed as a percentage of activity in the absence of \*\*\*RNA\*\*\*. Values shown are averages of duplicate reactions with standard deviations indicated. Background phosphorylation in the absence of DNA has been. . .

DETD The effect on DNA-PK activity was measured, relative to control reactions with no added \*\*\*RNA\*\*\*. Nonselected \*\*\*RNA\*\*\* had no effect on DNA-PK activity (FIG. 2A). Of the selected RNAs, seven inhibited DNA-PK activity by 85% or more. . .

DETD . . . in the presence of DNA. However, under these conditions, the DNA-PK was already highly active and a weak ability of \*\*\*RNA\*\*\* to activate DNA-PK might have gone undetected. Indeed, displacement of DNA by a weakly activating \*\*\*RNA\*\*\* may have resulted in partial inhibition of DNA-PK activity.

DETD . . . described above in the presence or absence of 0.25 nM double-stranded DNA, as indicated. Reactions contained various amounts of aptamer \*\*\*RNA\*\*\* (8, 16, 32 nM).

DETD When neither \*\*\*RNA\*\*\* nor DNA was present, DNA-PK had a low level of basal activity, as expected. The addition of double-stranded DNA increased. . .

DETD In principle, an \*\*\*RNA\*\*\* that binds to Ku protein could inhibit DNA-PK either by blocking the binding of Ku protein to DNA or by. . .

DETD . . . double stranded DNA (1 nM) oligonucleotide in the presence or absence of various amounts (8, 16, 32 nM) of nonradiolabeled \*\*\*RNA\*\*\*. The incubation conditions were 25 mM Tris-HCl pH 7.9, 0.5 mM EDTA, 10% glycerol, 5 mM MgCl<sub>2</sub>, 0.5 mM DTT, . . . inhibitors of DNA-PK enzyme activity also proved to be effective inhibitors of Ku protein-DNA binding activity (FIG. 2B). By contrast, \*\*\*RNA\*\*\* from the nonselected pool had no effect on Ku protein-DNA interaction (FIG. 2B). There was a rough correlation between the. . .

DETD . . . mapped, it is likely that it contains basic residues that are capable of electrostatic interactions with the phosphate backbone of \*\*\*RNA\*\*\* and DNA.

DETD To further explore the relationship between \*\*\*RNA\*\*\* and DNA binding sites in Ku protein, Electrophoretic Mobility Shift Assays (EMSAs) were performed comparing radiolabeled aptamer \*\*\*RNA\*\*\* (1 nM) and radiolabeled twenty-one base pair double stranded DNA (1 nM) probes. Cross-competition between \*\*\*RNA\*\*\* aptamers and DNA for Ku protein binding were found. It was discovered that Ku protein forms stable complexes with both types of probes, and that the complexes with \*\*\*RNA\*\*\* and DNA have similar electrophoretic mobilities. Both of the RNAs that were tested, #7-3 and #SC9, competed with DNA for binding to Ku protein. \*\*\*RNA\*\*\* #7-3 is a more effective competitor than \*\*\*RNA\*\*\* #SC9, consistent with the results of nitrocellulose filter binding assays. In the reciprocal experiment, nonradiolabeled DNA fragments competed with labeled \*\*\*RNA\*\*\* for binding to Ku protein. As controls, DNA and each \*\*\*RNA\*\*\* were shown to effectively self-compete for binding to Ku protein, confirming the fidelity of the analysis.

DETD Notably, neither the DNA nor the \*\*\*RNA\*\*\* competitors induced the formation of supershifted complexes. The absence of supershifted complexes provides additional evidence that \*\*\*RNA\*\*\* and DNA cannot bind simultaneously to Ku protein, but rather, bind competitively to the same site.

DETD . . . .mu.g protein) and other components as in FIG. 2. Reactions were performed in the presence of various amounts of aptamer \*\*\*RNA\*\*\* (250, 500, 1000 nM) as indicated. The assays were as described 20 above except a higher amount of DNA was. . . 100 to 142.1 .mu.M p53 was used. DNA-PK activity is expressed as a percentage of activity in the absence of \*\*\*RNA\*\*\*. Values shown are averages of duplicate reactions with standard deviations as indicated. Four different aptamer RNAs, all of which had. . . been shown to inhibit purified DNA-PK, were tested in the crude extract. All four reduced enzyme activity to near-basal levels. \*\*\*RNA\*\*\* from the nonselected pool had no effect at concentrations up to 1000 nM (FIG. 3, lanes 5-7). These results demonstrate. . .

DETD . . . (0.38 nM) under conditions for forming complexes for 30 minutes at room temperature. The complexes were challenged with either aptamer \*\*\*RNA\*\*\* or \*\*\*RNA\*\*\* from the nonselected pool. Incubation was then continued and the amount of Ku-DNA complexes remaining were measured by electrophoretic mobility. . .

DETD . . . the presence of aptamer #7-3. In contrast, the amount of Ku protein-DNA complexes remained almost unchanged when challenged with nonselected \*\*\*RNA\*\*\*. This result demonstrates that the aptamer \*\*\*RNA\*\*\* can compete with double-stranded DNA for the binding of Ku protein even if the aptamer \*\*\*RNA\*\*\* is added after Ku-DNA complexes have formed.

Set	Items	Description
S1	23	NUCLEOLAR (S) RNA (S) TAR
S2	13	RD (unique items)
S3	34	NUCLEOLAR AND TAR AND RNA
S4	22	RD (unique items)

>>>KWIC option is not available in file(s): 399

**4/3,K/1 (Item 1 from file: 5)**  
 DIALOG(R)File 5:Biosis Previews(R)  
 (c) 2003 BIOSIS. All rts. reserv.

14021055 BIOSIS NO.: 200300015084

**A \*nucleolar\* \*TAR\* decoy inhibitor of HIV-1 replication.**

AUTHOR: Michienzi Alessandro; Li Shirley; Zaia John A; Rossi John J(a)  
 AUTHOR ADDRESS: (a)Divisions of Molecular Biology, Beckman Research  
 Institute of the City of Hope, 1450 East Duarte Road, Duarte, CA,  
 91010-3011, USA\*\*USA E-Mail: jrossi@bricoh.edu

JOURNAL: Proceedings of the National Academy of Sciences of the United  
 States of America 99 (22):pl4047-14052 October 29 2002 2002

MEDIUM: print

ISSN: 0027-8424

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

**A \*nucleolar\* \*TAR\* decoy inhibitor of HIV-1 replication.**

...ABSTRACT: critical regulatory factor in HIV-1 gene expression. It mediates the transactivation of transcription from the HIV-1 LTR by binding to the transactivation response (\*TAR\*) element in a complex with cyclin T1. Because of its critical and early role in HIV gene expression, Tat and its interaction with the \*TAR\* element constitute important therapeutic targets for the treatment of HIV-1 infection. Based on the known \*nucleolar\* localization properties of Tat, we constructed a chimeric small \*nucleolar\* \*RNA\*-\*TAR\* decoy that localizes to the nucleoli of human cells and colocalizes in the nucleolus with a Tat-enhanced GFP fusion protein. When the chimeric \*RNA\* was stably expressed in human T lymphoblastoid CEM cells it potently inhibited HIV-1 replication. These results demonstrate that the \*nucleolar\* trafficking of Tat is critical for HIV-1 replication and suggests a role for the nucleolus in HIV-1 viral replication.

DESCRIPTORS:

...BIOSYSTEMATIC NAMES: DNA and \*RNA\* Reverse Transcribing Viruses,  
 Viruses, Microorganisms

**4/3,K/2 (Item 2 from file: 5)**  
 DIALOG(R)File 5:Biosis Previews(R)  
 (c) 2003 BIOSIS. All rts. reserv.

07779643 BIOSIS NO.: 000092083014

**HETEROLOGOUS BASIC DOMAIN SUBSTITUTIONS IN THE HIV-1 TAT PROTEIN REVEAL AN ARGININE-RICH MOTIF REQUIRED FOR TRANSACTIVATION**

AUTHOR: SUBRAMANIAN T; GOVINDARAJAN R; CHINNADURAI G

AUTHOR ADDRESS: INST. MOL. VIROL., ST. LOUIS UNIV. MED. CENT., 3681 PARK  
 AVE., ST. LOUIS, MO. 63110, USA.

JOURNAL: EMBO (EUR MOL BIOL ORGAN) J 10 (8). 1991. 2311-2318. 1991

FULL JOURNAL NAME: EMBO (European Molecular Biology Organization) Journal

CODEN: EMJOD

RECORD TYPE: Abstract

LANGUAGE: ENGLISH

...ABSTRACT: Tat protein coded by HIV-1 is a unique eukaryotic transactivator. It activates gene expression from the viral LTR by its interaction with a nascent \*RNA\* element (\*TAR\*) located at the 5' end of all HIV-1 transcripts. Tat appears to bind to its target \*RNA\* structure

in a highly sequence-specific manner. The \*TAR\*-binding activity of Tat has been localized in an Arg-rich basic domain located between residues 49 and 57 of the Tat protein. We have carried out domain substitution studies with heterologous basic domains which are also implicated in \*RNA\* binding. Here, we report that a 19 or a 12 amino acid region from the N-terminus of HTLV-I Rex can functionally substitute for...

...to the nucleus and nucleolus. The chimeric Tat proteins containing the basic domains of Rex and the N proteins are also localized in the nuclear/\*nucleolar\* region. Since the functionally inactive Tat - N chimeric proteins are still efficiently targeted to the nuclear/\*nucleolar\* compartments, our results suggest that nuclear/\*nucleolar\* localization alone is not sufficient for transactivation and demonstrate a direct role for the Arg motif in Tat function other than that required for nuclear/\*nucleolar\* localization.

4/3,K/3 (Item 3 from file: 5)  
DIALOG(R)File 5:Biosis Previews(R)  
(c) 2003 BIOSIS. All rts. reserv.

07320762 BIOSIS NO.: 000090100662

**A BULGE STRUCTURE IN HIV-1 \*TAR\* \*RNA\* IS REQUIRED FOR TAT BINDING AND TAT-MEDIATED TRANS-ACTIVATION**

AUTHOR: ROY S; DELLING U; CHEN C-H; ROSEN C A; SONENBERG N  
AUTHOR ADDRESS: DEP. BIOCHEM., MCGILL UNIV., MONTREAL, QUEBEC, H3G 1Y6, CANADA.

JOURNAL: GENES DEV 4 (8). 1990. 1365-1373. 1990

FULL JOURNAL NAME: Genes & Development

CODEN: GEDEE

RECORD TYPE: Abstract

LANGUAGE: ENGLISH

**A BULGE STRUCTURE IN HIV-1 \*TAR\* \*RNA\* IS REQUIRED FOR TAT BINDING AND TAT-MEDIATED TRANS-ACTIVATION**

...ABSTRACT: immunodeficiency virus type 1 (HIV-1) trans-activates viral gene expression and is obligatory for virus replication. Tat function is mediated through a sequence termed \*TAR\* that comprises part of the 5'-noncoding region of all HIV-1 mRNAs. This region forms a stable stem-loop structure in vitro. Recent evidence indicates that Tat binds directly to the \*TAR\* \*RNA\* sequence, and this binding is independent of the nucleotide sequence in the loop but dependent on the integrity of the upper stem. We used the...

...structure specificity of this interaction and its correlation with Tat trans-activation. We show that a 3-nucleotide bulge structure (positions +23 to +25) in \*TAR\* \*RNA\* is important for both Tat interaction with \*TAR\* \*RNA\* and Tat-mediated trans-activation of gene expression. Single base substitutions at position +23 that impair Tat-mediated trans-activation in vivo also reduce binding of Tat to \*TAR\* in vitro, suggesting that the first uridine residue in the bulge is the critical base for both functions. In contrast, mutations in the loop (positions...

...reduce Tat-mediated trans-activation, had no effect on Tat binding. We also show that a Tat peptide that includes the basic region required for \*nucleolar\* localization binds to \*TAR\* \*RNA\* with the same specificity as the full-length protein. We conclude that Tat binding to \*TAR\* is necessary but not sufficient by itself to account for trans-activation.

4/3,K/4 (Item 4 from file: 5)  
DIALOG(R)File 5:Biosis Previews(R)  
(c) 2003 BIOSIS. All rts. reserv.

07294844 BIOSIS NO.: 000090074731

**A TRANSDOMINANT TAT MUTANT THAT INHIBITS TAT-INDUCED GENE EXPRESSION FROM**



**THE HUMAN IMMUNODEFICIENCY VIRUS LONG TERMINAL REPEAT**

AUTHOR: PEARSON L; GARCIA J; WU F; MODESTI N; NELSON J; GAYNOR R  
AUTHOR ADDRESS: DIV. HEMATOL.-ONCOL., DEP. MED., UNIV. OF CALIFORNIA-LOS  
ANGELES, SCH. OF MED., LOS ANGELES, CALIF. 90024.  
JOURNAL: PROC NATL ACAD SCI U S A 87 (13). 1990. 5079-5083. 1990  
FULL JOURNAL NAME: Proceedings of the National Academy of Sciences of the  
United States of America  
CODEN: PNASA  
RECORD TYPE: Abstract  
LANGUAGE: ENGLISH

...ABSTRACT: virus (HIV) gene expression is dependent on specific  
regulatory regions in the long terminal repeat. These regions include the  
enhancer, SP1, "TATA," and trans-activating (\*TAR\*) regions. In addition,  
viral regulatory proteins such as tat and rev are important in regulating  
HIV gene expression. The mechanism of tat activation remains the...

...was localized predominantly to the nucleolus, .DELTA.tat was present in  
both the nucleus and cytoplasm, suggesting that it may inhibit tat  
function by preventing \*nucleolar\* localization. Transdominant mutants of  
tat may have a role in potentially inhibiting HIV gene expression.  
DESCRIPTORS: DNA MESSENGER \*RNA\* TAT PROTEIN

**4/3,K/5 (Item 1 from file: 34)**

DIALOG(R)File 34:SciSearch(R) Cited Ref Sci  
(c) 2003 Inst for Sci Info. All rts. reserv.

07360640 Genuine Article#: 155QX No. References: 81

**Title: The yeast *Saccharomyces cerevisiae* YDL112w ORF encodes the putative  
2'-O-ribose methyltransferase catalyzing the formation of Gm18 in tRNAs**

Author(s): Cavaille J; Chetouani F; Bachellerie JP (REPRINT)

Corporate Source: UNIV TOULOUSE 3,CNRS, LBME, 118 ROUTE NARBONNE/F-31062  
TOULOUSE//FRANCE/ (REPRINT); UNIV TOULOUSE 3,CNRS, LBME/F-31062  
TOULOUSE//FRANCE/

Journal: RNA-A PUBLICATION OF THE RNA SOCIETY, 1999, V5, N1 (JAN), P66-81

ISSN: 1355-8382 Publication date: 19990100

Publisher: CAMBRIDGE UNIV PRESS, 40 WEST 20TH STREET, NEW YORK, NY  
10011-4211

Language: English Document Type: ARTICLE (ABSTRACT AVAILABLE)

Abstract: The protein sequences of three known \*RNA\* 2'-O-ribose methylases  
were used as probes for detecting putative homologs through iterative  
searches of genomic databases. We have identified 45 new positive Open  
...

...organism, is considerably larger than its *Escherichia coli* functional  
homolog spoU (1,436 amino acids vs. 229 amino acids), or any known or  
putative prokaryotic \*RNA\* ribose methyltransferase. Homologs found in  
human (TRP-185 protein), *Caenorhabditis elegans* and *Arabidopsis*  
*thaliana* also exhibit a very long N-terminal extension not related to  
...

...Identifiers--SMALL \*NUCLEOLAR\* RNAs; ADENOSYLMETHIONINE-DEPENDENT  
METHYLTRANSFERASES; TRANSACTIVATING REGION \*RNA\*; HIV-1 \*TAR\* \*RNA\*;  
RIBOSOMAL-\*RNA\*; *ESCHERICHIA-COLI*; SEQUENCE MOTIFS; MESSENGER-\*RNA\*;  
SUBNUCLEAR LOCALIZATION; PSEUDOURIDINE SYNTHASE

**4/3,K/6 (Item 2 from file: 34)**

DIALOG(R)File 34:SciSearch(R) Cited Ref Sci  
(c) 2003 Inst for Sci Info. All rts. reserv.

06685289 Genuine Article#: ZK562 No. References: 35

**Title: \*Nucleolar\* protein p120 contains an arginine-rich domain that binds  
to ribosomal \*RNA\***

Author(s): Gustafson WC; Taylor CW; Valdez BC; Henning D; Phippard A; Ren Y  
; Busch H; Durban E (REPRINT)

Corporate Source: BAYLOR COLL MED, DEPT PHARMACOL, 1 BAYLOR  
PLAZA/HOUSTON//TX/77030 (REPRINT); BAYLOR COLL MED, DEPT  
PHARMACOL/HOUSTON//TX/77030  
Journal: BIOCHEMICAL JOURNAL, 1998, V331, 2 (APR 15), P387-393  
ISSN: 0264-6021 Publication date: 19980415  
Publisher: PORTLAND PRESS, 59 PORTLAND PLACE, LONDON, ENGLAND W1N 3AJ  
Language: English Document Type: ARTICLE (ABSTRACT AVAILABLE)

**Title: \*Nucleolar\* protein p120 contains an arginine-rich domain that binds to ribosomal \*RNA\***

...Abstract: Northwestern blotting procedure this study shows that recombinant p120 binds to an rRNA fragment in vitro with a dissociation constant of 4 nM. The specific \*RNA\*-binding region of p120 (residues 1-57) was identified with glutathione S-transferase-fused p120 deletion constructs and Northwestern blotting procedures. This \*RNA\*-binding region of p120, which includes the \*nucleolar\* localization signal of p120, is similar to the arginine-rich \*RNA\*-binding regions found in other \*RNA\*-binding proteins such as HIV Rev and Tat. Experiments in viuo with HeLa cell \*nucleolar\* extracts showed that p120 was associated with the 60-80 S pre-ribosomal particles. This association is disrupted by treatment with either RNase A or...

...Identifiers--POLYACRYLAMIDE-GEL ELECTROPHORESIS; HEPATOMA ASCITES-CELLS; RAT-LIVER; \*TAR\* \*RNA\*; PROLIFERATION; LOCALIZATION; RECOGNITION; NUCLEOPHOSMIN/B23; IDENTIFICATION; HOMOLOGY

4/3,K/7 (Item 3 from file: 34)

DIALOG(R)File 34:SciSearch(R) Cited Ref Sci  
(c) 2003 Inst for Sci Info. All rts. reserv.

05805295 Genuine Article#: WY721 No. References: 36

**Title: The post-transcriptional regulator Rev of HIV: Implications for its interaction with the \*nucleolar\* protein B23**

Author(s): Miyazaki Y (REPRINT) ; Nosaka T; Hatanaka M

Corporate Source: KYOTO UNIV, INST VIRUS RES, SAKYO KU/KYOTO 606//JAPAN/  
(REPRINT); SHIONOGI INST MED SCI, /OSAKA 566//JAPAN/

Journal: BIOCHIMIE, 1996, V78, N11-12, P1081-1086

ISSN: 0300-9084 Publication date: 19960000

Publisher: EDITIONS SCIENTIFIQUES MEDICALES ELSEVIER, 141 RUE JAVEL, 75747  
PARIS CEDEX 15, FRANCE

Language: English Document Type: ARTICLE (ABSTRACT AVAILABLE)

**Title: The post-transcriptional regulator Rev of HIV: Implications for its interaction with the \*nucleolar\* protein B23**

...Abstract: is controlled by trans-acting proteins. HIV-1 encodes several regulatory proteins, including two essential transactivations for viral replication, Rev and Tat. Both Rev and \*Tar\* have a \*nucleolar\* targeting signal and are actually located predominantly in the nucleoli. Within the nucleoli, Rev is localized to the combined regions of the dense fibrillar (DFC) and the granular (GC) components Tat does not colocalize precisely with any \*nucleolar\* component tested, but partly overlaps regions of the DFC and the GC. Regions of both Rev and Tat are overlapped by the distribution of the major \*nucleolar\* protein B23. Overexpression of Rev causes \*nucleolar\* ballooning and general structural deformity with aberrant accumulation of rRNAs, whereas Tat does not have that effect. B23 is markedly accumulated in those nucleoli deformed by Rev. Components of the \*nucleolar\* DFC, GC, and fibrillar center domains are not accumulated but dispersed in a few small spots or larger patches within the enlarged nucleoli.  
Cytophotometric DNA...

4/3,K/8 (Item 4 from file: 34)

DIALOG(R)File 34:SciSearch(R) Cited Ref Sci  
(c) 2003 Inst for Sci Info. All rts. reserv.

02573723 Genuine Article#: LM272 No. References: 38

**Title: \*TAR\* LOOP-DEPENDENT HUMAN-IMMUNODEFICIENCY-VIRUS TRANSACTIVATION  
REQUIRES FACTORS ENCODED ON HUMAN CHROMOSOME-12**

Author(s): HART CE; GALPHIN JC; WESTHAFFER MA; SCHUCHETMAN G

Corporate Source: CTR DIS CONTROL & PREVENT, DIV HIV AIDS, LAB INVEST BRANCH  
D12, 1600 CLIFTON RD/ATLANTA//GA/30333

Journal: JOURNAL OF VIROLOGY, 1993, V67, N8 (AUG), P5020-5024

ISSN: 0022-538X

Language: ENGLISH Document Type: NOTE (Abstract Available)

**Title: \*TAR\* LOOP-DEPENDENT HUMAN-IMMUNODEFICIENCY-VIRUS TRANSACTIVATION  
REQUIRES FACTORS ENCODED ON HUMAN CHROMOSOME-12**

Abstract: The trans-activator response region (\*TAR\*) \*RNA\* in the human immunodeficiency virus type 1 (HIV-1) and HIV-2 long terminal repeat forms stem-loop secondary structures in which the loop sequence is essential for trans activation. We investigated how the HIV trans-activation mechanism encoded on human chromosome 12 relates to the \*TAR\* \*RNA\* loop-dependent pathway. DNA transfection experiments showed that trans activation in human-hamster hybrid cells with the single human chromosome 12 and human T-cell lines was highly dependent on the native sequences of the HIV-1 \*TAR\* loop and the HIV-2 5' \*TAR\* loop. In nonhuman cell lines or hybrid cells without chromosome 12 that supported trans activation, the cellular mechanism was independent of the HIV-1 \*TAR\* loop and the response to mutations in the HIV-2 \*TAR\* loops differed from that found in human T-cell lines and human-hamster hybrid cells with chromosome 12. Our results suggest that the human chromosome 12 mechanism interacts directly with the \*TAR\* \*RNA\* loop or indirectly by regulating \*TAR\* \*RNA\*-binding proteins.

...Identifiers--MEDIATED TRANSACTIVATION; GENE-EXPRESSION; BINDING PROTEIN; NUCLEAR-PROTEIN; \*RNA\* SEQUENCES; RODENT CELLS; HIV-1 LTR; TYPE-1; REGION; ELEMENT

Research Fronts: 91-0190 001 (HUMAN-IMMUNODEFICIENCY-VIRUS TYPE-1; CHRONICALLY INFECTED CD8+ CELLS; COMPATIBILITY OF REV GENE ACTIVITY)

91-2472 001 (\*NUCLEOLAR\* ORGANIZER REGIONS; CELL CYCLE-DEPENDENT PHOSPHORYLATION OF HUMAN DNA POLYMERASE-ALPHA; ABERRANT KI-67 EXPRESSION)

91-3227 001 (\*RNA\* SECONDARY STRUCTURE; PRIMARY SEQUENCE; RHO-INDEPENDENT ESCHERICHIA-COLI TRANSCRIPTION TERMINATORS)

91-5983 001 (TRANSCRIBED REGION OF THE HUMAN C-FOS GENE; TISSUE-SPECIFIC ENHANCER ELEMENTS...

**4/3,K/9 (Item 5 from file: 34)**

DIALOG(R)File 34:SciSearch(R) Cited Ref Sci

(c) 2003 Inst for Sci Info. All rts. reserv.

01613878 Genuine Article#: HL816 No. References: 20

**Title: SPECIFIC BINDING OF ARGININE TO \*TAR\* \*RNA\***

Author(s): TAO JS; FRANKEL AD

Corporate Source: WHITEHEAD INST BIOMED RES, 9 CAMBRIDGE  
CTR/CAMBRIDGE//MA/02142

Journal: PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, 1992, V89, N7 (APR 1), P2723-2726

Language: ENGLISH Document Type: ARTICLE (Abstract Available)

**Title: SPECIFIC BINDING OF ARGININE TO \*TAR\* \*RNA\***

...Abstract: arginine residue within the basic region of the human immunodeficiency virus Tat protein mediates specific binding of Tat peptides to a three-nucleotide bulge in \*TAR\* \*RNA\*. It has been proposed that arginine recognizes \*TAR\* by forming a network of hydrogen bonds with two structurally distinct phosphates, an interaction termed the "arginine fork." Here it is shown that L-arginine blocks the Tat peptide/\*TAR\* interaction, whereas L-lysine and analogs of arginine that remove specific hydrogen bond donors do not. Experiments using an L-arginine affinity column demonstrate that arginine and the Tat peptides bind to the same site in \*TAR\*. Modification of two phosphates located at the junction of the double-stranded stem and bulge and modification of two adenine N7

groups in base-paired regions of \*TAR\* interfere with specific arginine binding. The results emphasize the importance of \*RNA\* structure in \*RNA\*-protein recognition and provide methods to identify arginine-binding sites in RNAs.

Research Fronts: 90-1803 001 (5S RIBOSOMAL-\*RNA\* SEQUENCES; U3 SMALL \*NUCLEOLAR\* RIBONUCLEOPROTEIN FUNCTIONS; BACTERIAL EVOLUTION)  
90-8119 001 (ESCHERICHIA-COLI METHIONYL-TRANSFER \*RNA\*-SYNTHETASE; MUTUALLY EXCLUSIVE SETS OF SEQUENCE MOTIFS; FRAMEWORK FOR UNDERSTANDING HOMEODOMAIN-DNA INTERACTIONS)

4/3,K/10 (Item 1 from file: 98)

DIALOG(R)File 98:General Sci Abs/Full-Text  
(c) 2003 The HW Wilson Co. All rts. reserv.

04517797 H.W. WILSON RECORD NUMBER: BGSA01017797 (USE FORMAT 7 FOR FULLTEXT)

#### Viruses and interferons.

Sen, Ganes C

Annual Review of Microbiology v. 55 (2001) p. 255-81

SPECIAL FEATURES: bibl il ISSN: 0066-4227

LANGUAGE: English

COUNTRY OF PUBLICATION: United States

WORD COUNT: 12173

(USE FORMAT 7 FOR FULLTEXT)

...ABSTRACT: cell signaling used by many cytokines and growth factors. Surprisingly, some of the same genes can also be induced directly by viruses and double-stranded \*RNA\*, a common viral by-product. Some of the interferon-induced proteins have novel biochemical properties and some are inactive as such but can be activated by double-stranded \*RNA\* produced during virus infection. Finally, almost all viruses have evolved mechanisms to evade the interferon system by partially blocking interferon synthesis or interferon action. Thus...

#### TEXT:

Key Words antiviral effects, double-stranded \*RNA\*, interferon-induced proteins, transcriptional regulation, Jak-STAT pathway

#### INTRODUCTION TO THE INTERFERON SYSTEM

Interferons (IFNs) are the first known members of the important biological regulatory...

...IFN-induced proteins remain unknown, some of the proteins that have been analyzed carry out interesting functions. Moreover, two such proteins use double-stranded (ds) \*RNA\* as a cofactor for their enzymatic activity, providing the first biochemical function of dsRNA.

In the context of host-virus infection, regulation of components of... activation of IFN-a/b genes is mediated by complex pathways involving many transcription factors. A well-characterized pathway is triggered by viral double-stranded \*RNA\* expressed in infected cells. Other lesser-known pathways that are activated by viral interactions with cellular receptors or by viral proteins must exist as well...

...the cytoplasm and are activated by phosphorylation. NFkB activation involves activation of the Ikb kinase complex and the dsRNA-dependent protein kinase, PKR (protein kinase, \*RNA\*-activated) (12). P38 and JNK (Jun N-terminal kinase) kinases are also activated by MEKK1 (microtubule-associated protein, extracellular response kinase kinase) (39). Complete characterization...length of double-stranded region is good enough for activating them maximally (72). Consequently, many viral RNAs with partial ds structures, such as adenoviral VAI \*RNA\*, EB viral EBER \*RNA\*, and HIV-1 \*TAR\* \*RNA\*, can activate them (74). The small enzymes are tetramers and the medium isozymes are dimers; their oligomerization is needed for their enzyme activity. The catalytic...c-Myc, and Rb (15). The

inhibition of c-Myc activity is mediated by blocking its association with Max (94). Another member, P204, is a \*nucleolar\* protein that inhibits ribosomal \*RNA\* transcription by binding to an essential transcription factor, UBF-1 (50). There are indications that these proteins may have cellular functions beyond the IFN system...EMCV) (26). This effect requires enzymatically active proteins and the presence of RNaseL, indicating the need for 2-5(A) production and activated RNaseL-mediated \*RNA\* degradation (33). Viral dsRNA is the likely activator of 2-5(A) synthetases because they coimmunoprecipitate to form an enzymatically active complex (28). Although the...signaling are HBV terminal protein, EBV EBNA1 and EBNA2 proteins, HPV E7 protein, polyoma virus large T, and HCMV and HHV8 proteins (7, 27). Numerous \*RNA\* viruses also interfere with IFN signaling. These include influenza virus, ebola virus, Sendai virus, SV5, bovine respiratory syncytial virus (BRSV), parainfluenza virus, and hepatitis C...encode RNAs that have partial double-stranded structures, and at high concentrations they block PKR activation by authentic dsRNA. Such viral RNAs include adenoviral VAI \*RNA\*, EBV EBER \*RNA\*, and HIV-1 \*TAR\* \*RNA\* (7, 27). Vaccinia virus E3L protein, influenza virus NS1 protein (5), and reovirus sigma 3 protein bind dsRNA and thus can block PKR activation. The...

...through unknown mechanisms. HSV1 encodes two proteins that can interfere with PKR activation or action (86). One of these proteins is Us11, a virally encoded \*RNA\*-binding protein (9). The other is the g1 34.5 protein. The carboxyl terminus of this protein binds to the cellular protein phosphatase 1a and...

...antiproliferative activity of IFN- $\alpha$ . Differential IFN- $\alpha$  sensitivity of different clonal derivatives of Daudi cells has been correlated with the abundance of a viral \*RNA\* that has considerable double-stranded structure (24). This \*RNA\* may activate PKR, leading to translation inhibition, c-Myc repression, and NFkB activation, causing cell growth inhibition. IFN- $\gamma$  may promote cell proliferation or apoptosis...grants from the National Institutes of Health.

Figure 1 Interactions between viruses and the interferon system. Viruses enter the cell, get uncoated, and synthesize viral \*RNA\* and proteins. Often viral dsRNA is also produced. dsRNA can signal to the promoter of interferon (IFN)- $\alpha$ /b genes by activating the transcription factors...

...to viral infection. Immunity 13:129-41

4. Bandyopadhyay SK, Leonard GT Jr, Bandyopadhyay T, Stark GR, Sen GC. 1995. Transcriptional induction by double-stranded \*RNA\* is mediated by interferon-stimulated response elements without activation of interferon-stimulated gene factor 3. J. Biol. Chem. 270:19624-29

5. Bergmann M, Garcia...and IKKbeta are required for activating the innate response to viral infection. Immunity 11:721-31

13. Clemens MJ, Elia A. 1997. The double-stranded \*RNA\*-dependent protein kinase PKR: structure and function. J. Interferon Cytokine Res. 17:503-24

14. Darnell JE Jr, Kerr IM, Stark GR. 1994. Jak-STAT...

...virus countermeasures. J. Gen. Virol. 81:2341-64

28. Griboaud G, Lembo D, Cavallo G, Landolfo S, Lengyel P. 1991. Interferon action: binding of viral \*RNA\* to the 40-kilodalton 2'-5'-oligoadenylate synthetase in interferon-treated HeLa cells infected with encephalomyocarditis virus. J. Virol. 65:1748-57

29. Guo J...

...factor 3. EMBO J. 19:6891-99

30. Guo J, Peters K, Sen GC. 2000. Induction of the human protein P56 by interferon, double-stranded \*RNA\* or virus infection. Virology 267:209-19

31. Guo J, Sen GC. 2000. Characterization of the interaction between the interferon-induced protein P56 and the...

...the mouse mammary tumor virus. J. Virol. 74:1892-99

32. Haller O, Frese M, Kochs G. 1998. Mx proteins: mediators of innate resistance to \*RNA\* viruses. Rev. Sci. Tech. 17:220-30

33. Hassel BA, Zhou A, Sotomayor C, Maran A, Silverman RH. 1993. A dominant negative mutant of 2...

...IFN-alpha, but is required by the mouse cytomegalovirus for its replication. Virology 262:1-8

36. Hovanessian AG. 1991. Interferon-induced and double-stranded \*RNA\*-activated enzymes: a specific protein kinase and 2',5'-oligoadenylate synthetases. J. Interferon Res. 11:199-205

37. Ito T, Yang M, May WS. 1999. RAX, a cellular activator for double-stranded \*RNA\*-dependent protein kinase during stress signaling. J. Biol. Chem. 274:15427-32

38. Janzen C, Kochs G, Haller O. 2000. A monomeric GTPase-negative MxA ...17:738-50

43. Leaman DW, Salvekar A, Patel R, Sen GC, Stark GR. 1998. A mutant cell line defective in response to double-stranded \*RNA\* and in regulating basal expression of interferon-stimulated genes. Proc. Natl. Acad. Sci. USA 95:9442-47

44. Leib DA, Machalek MA, Williams BR, Silverman...

...of p204, a protein also inducible by interferon. Mol. Cell Biol. 20:7024-36

50. Liu CJ, Wang H, Lengyel P. 1999. The interferon-inducible \*nucleolar\* p204 protein binds the ribosomal \*RNA\*-specific UBF1 transcription factor and inhibits ribosomal \*RNA\* transcription. EMBO J. 18:2845-54

51. Lopez S, Island ML, Drouin J, Bandu MT, Christeff N, et al. 2000. Repression of virus-induced interferon...

...protein kinase, PKR. EMBO J. 17:4379-90

62. Patel RC, Stanton P, McMillan NM, Williams BR, Sen GC. 1995. The interferon-inducible double-stranded \*RNA\*-activated protein kinase self-associates in vitro and in vivo. Proc. Natl. Acad. Sci. USA 92:8283-87

63. Patel RC, Vestal DJ, Xu Z, Bandyopadhyay S, Guo W, et al. 1999. DRBP76, a double-stranded \*RNA\*-binding nuclear protein, is phosphorylated by the interferon-induced protein kinase, PKR. J. Biol. Chem. 274:20432-37

64. Peters G, Hartmann R, Qin J...interferons. Annu. Rev. Biochem. 67:227-64

80. Stojdl DF, Abraham N, Knowles S, Marius R, Brasey A, et al. 2000. The murine double-stranded \*RNA\*-dependent protein kinase PKR is required for resistance to vesicular stomatitis virus. J. Virol. 74:9580-85

81. Symons JA, Alcamí A, Smith GL. 1995...

...Tiwari RK, Kusari J, Sen GC. 1987. Functional equivalents of interferon-mediated signals needed for induction of an mRNA can be generated by double-stranded \*RNA\* and growth factors. EMBO J. 6:3373-78

90. Tortorella D, Gewurz BE, Furman MH, Schust DJ, Ploegh HL. 2000. Viral subversion of the immune...

...1:507-18

97. Weaver BK, Kumar KP, Reich NC. 1998. Interferon regulatory factor 3 and CREB-binding protein/p300 are subunits of double-stranded \*RNA\*-activated transcription factor DRAFI. Mol. Cell Biol. 18:1359-68

98. Whitten TM, Quets AT, Schloemer RH. 1991. Identification of the hepatitis B virus factor...

...Biochem. 70:489-506

101. Xu Z, Williams BR. 2000. The B56alpha regulatory subunit of protein phosphatase 2A is a target for regulation by doublestranded \*RNA\*-dependent protein kinase PKR. Mol. Cell Biol. 20:5285-99

102. Yie J, Senger K, Thanos D. 1999. Mechanism by which the IFN-beta enhanceosome...

04255630 H.W. WILSON RECORD NUMBER: BGSA00005630 (USE FORMAT 7 FOR FULLTEXT)

**Lentivirus replication and regulation.**

AUGMENTED TITLE: review

Tang, Hengli

Kuhen, Kelli L; Wong-Staal, Flossie

Annual Review of Genetics v. 33 (1999) p. 133-70

SPECIAL FEATURES: bibl il ISSN: 0066-4197

LANGUAGE: English

COUNTRY OF PUBLICATION: United States

WORD COUNT: 17942

(USE FORMAT 7 FOR FULLTEXT)

TEXT:

... immunodeficiency virus, transcriptional regulation,  
posttranscriptional control, downregulation

INTRODUCTION

GENOMIC STRUCTURES OF LENTIVIRUSES

Lentiviruses comprise one of three subfamilies of retroviruses, which also include the oncoviruses (\*RNA\* tumor viruses) and spuma (or foamy) viruses. Members of all three subfamilies are able to infect humans, and are associated with a spectrum of pathogenic...

...readers to the excellent recent text on Retroviruses (42). Rather, we give a broad outline here for the purpose of comparison with the lentiviruses. The \*RNA\* genome of retroviruses is flanked by two short, redundant (R) sequences at both termini. These are adjacent to unique sequences, U5 and U3, found at the 5' and 3' ends, respectively. The defining feature of a retrovirus is its ability to convert its \*RNA\* genome to a DNA intermediate through the virally encoded reverse transcriptase. A subsequent reaction catalyzed by another viral enzyme, integrase, results in the integration of...

...coupled and only completely spliced mRNA species are exported to the cytoplasm. However, retroviruses need to bypass this regulation in order to utilize unspliced viral \*RNA\* both as messenger \*RNA\* for protein synthesis and as genomic \*RNA\* for progeny virions. Interestingly, lentiviruses and simpler retroviruses utilize different strategies to this same end, as detailed in a subsequent section.

The virion capsid contains a diploid genome of two (+) \*RNA\* strands. A small region, usually spanning the 3' end of the LTR and the 5' end of the gag gene, mediates specific binding to the...a natural host cofactor for HIV-1 integration (33).

TRANSCRIPTIONAL REGULATION

Regulation of HIV-1 transcription involves a complex interplay between cis-acting DNA and \*RNA\* elements present within the chromatin-associated proviral long terminal repeats (LTRs), cellular transcription factors (DNA level), and the viral trans-regulatory protein Tat (\*RNA\* level).

The HIV-1 LTR and Regulatory Elements The long terminal repeat (LTR) DNA, defined by the U3, R, and U5 regions, possesses multiple cis...

...3). The LTR can be divided into several discrete functional domains: the negative regulatory element (NRE), the enhancer, the basal promoter elements, the core promoter, \*TAR\* (Tat-activation response element), and the IST (inducer of short transcripts). Of note are two unique cis-acting elements: IST, a DNA element, functions as an enhancer that promotes the synthesis of abortive transcripts; and \*TAR\*, and \*RNA\* element present in the viral LTR and the 5' end of all viral mRNAs, is an enhancer that promotes the synthesis of processive transcripts.

Upstream...

...impacts transcription, as the loss of this region results in an increase in gene expression. This region has a negative effect on the rate of \*RNA\* initiation in vivo (145) and in vitro (181). The NRE has numerous candidate protein-binding sites and many of these are occupied, as observed by...boxes may regulate a switch from basal to activated gene expression (186). The TATA box functions to recruit a coordinated series of transcription factors and \*RNA\* polymerase II. Initiation of transcription is preceded by binding of TFIID (transcription factor IID), a pivotal factor in the basal transcription apparatus. This is followed by assembly of \*RNA\* polymerase II and the general initiation factors with the core promoter to form the preinitiation complex (27, 223). The formation of this complex is sufficient...

...the action of Tat in enhancing transcription (187). There is also a candidate binding site for CTF (CAAT-box transcription factor) in this region.

TAT/\*TAR\* Interaction Tat is a potent transactivator of gene expression for HIV-1. The most interesting and unique function of Tat is that it acts through an \*RNA\* target termed \*TAR\* located immediately 3' to the LTR transcription start site. The 59-nt \*TAR\* element forms a stable, partially base-paired stem-loop structure. This stable secondary structure has been confirmed by in vitro chemical and enzyme mapping and by NMR analysis (120). There are two critical sequence elements in \*TAR\* that are required for tat transactivation: a hexanucleotide loop at the tip and a three-nucleotide uracil-rich bulge within the stem (Figure 4). The...

...Tat span the bulge and adjacent sequences (37). Although not essential for binding of Tat in vitro, the loop is also important for the Tat-\*TAR\* interaction in vivo since any mutation in the loop sequence would abolish such interactions (70, 211).

Tat is a 14-kDa protein comprised of 86...

...protein possesses the activation domain. Residues 48 through 57 comprise an arg/lys-rich basic region that is responsible both for direct interaction with the \*TAR\* \*RNA\* target and for nuclear/\*nucleolar\* localization. The arginine at aa position 52 is critical for this specific association with \*TAR\*. The carboxy-terminal region, encoded by exon 2, is dispensable for function.

At the transcriptional level, the Tat/\*TAR\* interaction can potentially enhance or stabilize binding of transcription complexes required for both initiation and elongation of viral RNAs (134, 154). However, now the general consensus is that the primary function of Tat is to increase the processivity of \*RNA\* polymerase II rather than stimulating transcription (205, 272). The increased initiation of transcription that is observed is probably a result of the higher rate of promoter...

...is a direct consequence of enhanced processivity. Much effort has been placed on identifying cellular cofactors of Tat. A number of cellular factors bind to \*TAR\*, but no clear role for any of these factors has been established in Tat regulation. Recently, a multisubunit cellular protein kinase called TAK (Tat-associated...

...was identified. TAK directly binds to the activation domain of Tat and, in turn, hyperphosphorylates the carboxyl-terminal domain (CTD) of the large subunit of \*RNA\* polymerase II (Pol II) (110, 267). The CTD possesses a heptad repeat consisting of the consensus sequence Tyr-Ser-Pro-Thr-Ser-Pro-Ser. Hypophosphorylated...

...of the TAK complex, directly binds to the activation domain of Tat (255). Furthermore, a functional complex requires the simultaneous binding of CycT1 to the \*TAR\* loop and Tat to the \*TAR\* bulge. CycT1 increases both the affinity and specificity of Tat for \*TAR\* (94). CycT1 was also identified as a regulatory subunit of Cdk9. As shown in Figure 4, the complex formation of Tat-CycT1 with \*TAR\* recruits CDK9 to the pol II transcription complex, ...factor encoded on human chromosome 12 (5, 105, 106). The observation that human cyclin T1 is encoded on chromosome 12,



enhances binding of Tat to \*TAR\* in vitro, and augments Tat transactivation in murine cells (255) suggests that the murine cyclin T protein is unable to support efficient Tat activation of...

...cysteine for tyrosine at position 261 in murine CycT restores zinc-dependent binding to Tat as well as high-affinity binding of the complex to \*TAR\* \*RNA\*. This single change was also found to be both necessary and sufficient for murine CycT to support high levels of Tat trans-activation in murine...

...stable expression of human CycT1 in murine cells was insufficient to overcome additional species-specific barriers to HIV-1 replication in murine cells.

Effects of \*TAR\* \*RNA\* and TAT on Translation The level of \*TAR\* \*RNA\* synthesized may activate or inhibit cellular repressors in different phases of virus replication. For example, high levels of \*TAR\*-containing \*RNA\* inhibit the interferon (IFN)-inducible, \*RNA\*-dependent protein kinase PKR (103). PKR is a pivotal mediator of the antiviral effects of IFN (216), and acts through phosphorylation of the  $\alpha$  subunit of the translation factor eIF-2 (217). Many viruses have evolved strategies to inhibit PKR function (126). Thus, inhibition of PKR by \*TAR\* \*RNA\* would allow viral protein synthesis and a productive infection to proceed. In contrast, low levels of \*TAR\* \*RNA\* activate PKR (210), which may help to maintain the latent phase of virus infection in the host cell. In addition to \*TAR\*-containing RNAs, the Tat protein can also inhibit PKR kinase activity by competing with eIF-2 $\alpha$  for binding to PKR (20, 162). Thus, it is conceivable that Tat and its viral \*RNA\* target sequence, \*TAR\*, can stimulate gene expression at the level of translation by inhibition of PKR function.

Downstream DNA Regulatory Elements In vitro and in vivo footprinting analyses...

...primary transcript upon transcription from the integrated proviral DNA, which serves as template for synthesis of gag and pol gene products as well as genomic \*RNA\* for encapsidation into progeny virions. It also serves as the precursor mRNA for further processing to generate multiple spliced mRNAs (244). Thus, retroviruses defy the...

...production. In addition to mediating nuclear export of unspliced mRNA, Rev has also been proposed to function through inhibition of splicing of the full-length \*RNA\* (32, 131) or increasing the efficiency of translation of the late gene products (6).

REV-RRE Interaction Rev of HIV-1 is a 116-amino activation domain. The N-terminal half of Rev contains sequences that are important for nuclear localization, multimerization, and \*RNA\*-binding. The Rev Response Element (RRE) binding site of Rev overlaps with the nuclear localization signal (NLS). Unlike the classical lysine-rich NLS, the Rev...

...b for nuclear import (109, 242). The activation domain at the carboxyl terminus of Rev contains a nuclear export signal (NES) that accesses a cellular \*RNA\* export pathway different from that of cellular mRNA (72). Mutations in this domain abolish the ability of Rev to shuttle between the cytoplasm and the...

...phenotype and are capable of repressing wild-type Rev function as well as HIV replication (149).

The RRE of HIV-1 is a highly structured \*RNA\* element that is present in the singly spliced env mRNA and the full-length genomic \*RNA\*. It serves as a high-affinity binding site of Rev on these \*RNA\* (150). Rev interacts with RRE in a multimerized form, directing RRE-containing \*RNA\* to a nuclear export pathway specified by its leucine-rich NES (activation domain). The Rev-RRE interaction has been studied extensively, both in vitro and in vivo. Rev binds to in vitro transcribed RRE \*RNA\* with high affinity and specificity. RRE contains extensive secondary structure, including several stem loops. The second stem from the 5' end, SLII, is the initial and primary binding site for Rev (11, 13). When an antibody directed against Rev was used to immunoprecipitate \*RNA\* from Rev and RRE

transfected cells, RRE-containing \*RNA\* was specifically detected, demonstrating the in vivo interaction between Rev and RRE (7, 28). Note that the RRE-binding domain of Rev overlaps with the...

...remains accessible on the RRE-bound Rev protein. Furthermore, RRE binding effectively masks the NLS of Rev, providing a mechanism to keep the RRE-containing \*RNA\* from being brought back to the nucleus by Rev (109).

Cellular Cofactors for REV/RRE Several cellular cofactors have been implicated in Rev-mediated \*RNA\* export. The cellular protein eIF-5A was shown to cross-link to Rev/RRE specifically in vitro, and a transdominant negative form of eIF-5A...

...and 5S rRNA use the same export pathway (72), eIF-5A may act as an adapter molecule between the RRE/Rev complex and a cellular \*RNA\* export pathway. Human RIP/Rab, a nucleoporin-like protein, was identified to interact with the activation domain of Rev by yeast two-hybrid screens and ...

...strain (173). The CRM-1/NES interaction is RanGTP dependent and sensitive to leptomycin B, a drug that blocks both Rev- and Rev/RRE-mediated \*RNA\* export (261).

Another positive cofactor in the Rev transactivation pathway is \*RNA\* helicase A (RHA). RHA was originally identified as a protein binding to the constitutive transport element (CTE) of type D retroviruses (see below) (239). We...

...is insensitive to leptomycin B (240). Therefore, we propose that RHA acts at a step prior to nuclear export in the Rev transactivation pathway. Since \*RNA\* helicases have been implicated in the release of mature mRNA from the spliceosomes (180, 182), a possible mechanism of RHA is in the premature release...

...140).

RRE-binding cellular proteins have also been identified. A novel nuclear factor with homology to heterogeneous nuclear ribonucleoprotein particle (hnRNP) F interacts with RRE \*RNA\* in vitro and may negatively regulate RRE-mediated gene expression (265). The generic splicing factor ASF/SF2 binds to RRE in a Rev-dependent manner...

...require the nuclear export of unspliced mRNA for the synthesis of viral structural proteins and genomic encapsidation. Instead of encoding transactivating proteins and their cognate \*RNA\* response elements, simple retroviruses encode cis-acting \*RNA\* elements that interact directly with the cellular \*RNA\* export ... exports via the 5S rRNA export pathway, CTE seems to utilize a distinct mRNA export pathway (72, 189, 214). More specifically, nuclear export of CTE \*RNA\* is independent of CRM-1 and is insensitive to leptomycin B inhibition (184).

Since CTE functions in the absence of any viral protein, it is likely that it interacts directly with components of a particular cellular export machinery by means of \*RNA\*-protein interaction. Two CTE-binding proteins, RHA and TAP, were recently identified (101, 239). As noted above, RHA contains a nuclear export signal that is insensitive to leptomycin B. However, whether RHA mediates the nuclear export of CTE \*RNA\* and its potential interaction with TAP remain to be determined.

A direct repeat (DR) sequence located between the env gene and the 3' LTR of...

...of the HIV Gag protein. RSV mutants that lack this element did not replicate well and seemed to have a significant reduction of unspliced viral \*RNA\* (179). Like MPMV and SRV-1, RSV does not encode a Rev-like protein, so it is likely that this cis-acting DR element also...

...promotes expression of CAT gene placed in an intron (57, 114, 116). However, the mechanism by which PRE functions to facilitate cytoplasmic appearance of unspliced \*RNA\* may not be restricted to direct \*RNA\* export (T Hope, personal communication). A 119-nucleotide \*RNA\* element within the herpes simplex virus thymidine kinase gene (a naturally intronless viral

gene) can also promote cytoplasmic accumulation of unspliced b-globulin transcript; this...

...binds hnRNP L (143). Additional cis-acting elements similar to PPE were also identified within the TK gene (185). Table 1 summarizes the various viral \*RNA\* elements for posttranscriptional regulation.

#### LENTIVIRUS-HOST INTERACTIONS

Infection by lentiviruses or expression of specific lentiviral gene products can have profound effects on phenotypes of the...entry has greatly propelled this area of research. Similarly, studies on the mechanism of Rev function have unraveled novel pathways for the nucleocytoplasmic transport of \*RNA\* and proteins. A more direct application of lentivirus research relates to the design of gene transfer vectors. The unique ability of lentiviruses to infect nondividing...  
Biology, University of California, San Diego, California 92093-0665;  
E-mail: htang@biomail.ucsd.edu; KKuhen@aol.com; fwongstaal@ucsd.edu

TABLE 1 Cis-acting \*RNA\* elements for posttranscriptional regulation of viral gene expression

Figure 1 Genomic structure of lentiviruses. The genomic organization of several lentiviruses is depicted. Moloney leukemia virus...

...regulatory factor binding site. (Adapted from 129,246.)

Figure 4 Schematic representation of the interaction of the Tat:Cyclin T:CDK9 heterotrimeric complex with the \*TAR\* \*RNA\* element. Tat is shown interacting with the bulge region, whereas cyclin T is shown interacting primarily with the loop region. (Adapted from 50.)

Figure 5 A model for the posttranscriptional regulation of HIV-1 \*RNA\* . \*RNA\* helicase A (RHA) is recruited to the viral mRNA splicing complex by directly binding to RRE and mediates the premature release of unspliced or incompletely...

...tropic HIV-1. Science 272:1955-58

5. Alonso A, Cujec T, Peterlin BM. 1994. Effects of human chromosome 12 on interactions between Tat and \*TAR\* of human immunodeficiency virus type 1. J. Virol. 68:6505-13

6. Arrigo SJ, Chen IS. 1991. Rev is necessary for translation but not cytoplasmic...

...94

11. Bartel DP, Zapp ML, Green MR, Szostak JW. 1991. HIV-1 Rev regulation involves recognition of non-Watson-Crick base pairs in viral \*RNA\*. Cell 67:529-36

12. Bartz SR, Rogel ME, Emerman M. 1996. Human immunodeficiency virus type 1 cell cycle control: Vpr is cytostatic and mediates...

...J. Virol. 70:2324-31

13. Battiste JL, Tan R, Frankel AD, Williamson JR. 1994. Binding of an HIV Rev peptide to Rev responsive element \*RNA\* induces formation of purine-purine base pairs. Biochemistry 33:2741-47

14. Berger EA. 1997. HIV entry and tropism: the chemokine receptor connection. AIDS 11...Proc. Natl. Acad. Sci. USA 89:6580-84

27. Buratowski S, Hahn S, Guarente L, Sharp PA. 1989. Five intermediate complexes in transcription initiation by \*RNA\* polymerase II. Cell 56:549-61

28. Campbell LH, Borg KT, Haines JK, Moon RT, Schoenberg DR, et al. 1994. Human immunodeficiency virus type 1...

...type 1 Tat. J. Virol. 72:2615-29

37. Churcher MJ, Lamont C, Hamy F, Ding-wall C, Green S. 1993. High affinity binding of \*TAR\* \*RNA\* by the human immunodeficiency virus Type-1 tat protein requires base-pairs in the \*RNA\* stem and amino acid residues flanking the basic region. J. Mol. Biol. 230:90-110

38. Clapham PR. 1997. HIV and chemokines: ligands sharing cell...Cell 87:745-56

66. Ernst RK, Bray M, Rekosh D, Hammarskjöld ML. 1997. Secondary structure and mutational analysis of the Mason-Pfizer monkey virus \*RNA\*

constitutive transport element. \*RNA\* 3:210-22

67. Ernst RK, Bray M, Rekosh D, Hammariskjold ML. 1997. A structured retroviral \*RNA\* element that mediates nucleocytoplasmic export of intron-containing \*RNA\*. Mol. Cell. Biol. 17:135-44

68. Esposito D, Craigie R. 1998. Sequence specificity of viral end DNA binding by HIV-1 integrase reveals critical...

...association between gp120 and CCR5. J. Virol. 72:1160-64

70. Feng S, Holland EC. 1988. HIV-1 tat transactivation requires the loop sequences within \*TAR\*. Nature 334:165-67

71. Feng Y, Broder CC, Kennedy PE, Berger EA. 1996. HIV-1 entry cofactor: functional cDNA cloning of a seven-transmembrane...89

101. Gruter P, Tabernero C, von Kobbe C, Schmitt C, Saavedra C, et al. 1998. TAP the human homolog of Mex67p mediates CTE-dependent \*RNA\* export from the nucleus. Mol. Cell 1:649-59

102. Guermah M, Malik S, Roeder RG. 1998. Involvement of TFIID and USA components in transcriptional...

...promoter by NF-kappaB and Spl. Mol. Cell. Biol. 18:3234-44

103. Gunnery S, Rice AP, Robertson HD, Mathews MB. 1990.

Tat-responsive region \*RNA\* of human immunodeficiency virus 1 can prevent activation of the double-stranded-\*RNA\*-activated protein kinase. Proc.

Natl. Acad. Sci. USA 87:8687-91

104. Guo B, Odgren PR, Wijnen VAJ, Last TJ, Nickerson J, et al. 1995

...

...NMP-1 is the transcription factor YY1. Proc. Natl. Acad. Sci. USA 92:10526-30

105. Hart C, Galphin J, Westhafter M, Schochetman G. 1993. \*TAR\* loop-dependent human immunodeficiency virus trans activation requires factors encoded on human chromosome 12. J. Virol. 67:5020-24

106. Hart C, Ou C, Galphin...

...Rice A. 1995. Lentivirus Tat proteins specifically associate with a cellular protein kinase, TAK, that hyperphosphorylates the carboxyl-terminal domain of the large subunit of \*RNA\* polymerase II: candidate for a Tat cofactor. J. Virol. 69:1612-20

111. Hill CM, Littman DR. 1996. Natural resistance to HIV? Nature 382:668...infectivity by MAPK, a virion-associated kinase. EMBO J. 17:2607-18

120. Jaeger JA, Tinoco I. 1993. An NMR study of the HIV-1 \*TAR\* element hairpin. Biochemistry 32:12522-30

121. Jenkins Y, McEntee M, Weis K, Greene WC. 1998. Characterization of HIV-1 vpr nuclear import: analysis of...

...protein. Science 268:429-31

140. Li J, Tang H, Mullen TM, Westberg C, Reddy TR, Rose DW, Wong-Staal F. 1999. A role for \*RNA\* helicase A in posttranscriptional regulation of HIV type 1. Proc. Natl. Acad. Sci. USA 96:709-14

141. Liang C, Li X, Quan Y, Laughrea...

...of some multiply-exposed individuals to HIV-1 infection. Cell 86:367-77

143. Liu X, Mertz JE. 1995. HnRNP L binds a cis-acting \*RNA\* sequence element that enables intron-dependent gene expression. Genes Dev. 9:1766-80

144. Liu YZ, Latchman DS. 1997. The octamer-binding proteins Oct-1...  
162. McMillan NA, Chun RF, Siderovsky DP, Galabru J, Toone WM, et al. 1995. HIV-1 Tat directly interacts with the interferon-induced, double-stranded \*RNA\*-dependent kinase, PKR. Virology 213:413-24

163. Michael NL, Chang G, Louie LG, Mascola JR, Dondero D, et al. 1997. The role of viral...

...be determined by regions of gp120 outside the CD4-binding domain. Nature 348:69-73

179. Ogert RA, Lee LH, Beemon KL. 1996. Avian retroviral \*RNA\* element promotes unspliced \*RNA\* accumulation in the cytoplasm. J. Virol. 70:3834-43

180. Ohno M, Shimura Y. 1996. A human \*RNA\* helicase-like protein HRH1

facilitates nuclear export of spliced mRNA by releasing the \*RNA\* from the spliceosome. *Genes Dev.* 10:997-1007

181. Okamoto T, Bonter T, Josephs SF, Sadaie MR, Wong-Staal F. 1990. Transcriptional activation from the long-terminal repeat of human immunodeficiency virus in vitro. *Virology* 177:606-14

182. Ono Y, Ohno M, Shimura Y. 1994. Identification of a putative \*RNA\* helicase (HRH1) a human homolog of yeast Prp22. *Mol. ...* 185. Otero GC, Hope TJ. 1998. Splicing-independent expression of the herpes simplex virus type 1 thymidine kinase gene is mediated by three cis-acting \*RNA\* subelements. *J. Virol.* 72:9889-96

186. Ou SH, Garcia-Martinez LF, Paulssen EJ, Gaynor RB. 1994. Role of flanking E box motifs in human...

...Maniatis T, Greene WC. 1997. HIV Rev-dependent binding of SF2/ASF to the Rev response element: possible role in Rev-mediated inhibition of HIV \*RNA\* splicing. *Proc. Natl Acad. Sci. USA* 94:973-78

203. Re F, Braaten D, Franke EK, Luban J. 1995. Human immunodeficiency virus type 1 Vpr...

...DNA depends on mitosis. *EMBO J.* 12:2099-108

208. Roebuck KA, Brenner DA, Kagnoff MF. 1993. Identification of c-fos-responsive elements downstream of \*TAR\* in the long terminal repeat of human immunodeficiency virus type-1. *J. Clin. Invest.* 92:1336-49

209. Roy AL, Meisterernst M, Pognonec P, Roeder...

...S, Agy M, Hovanessian AG, Sonenberg N, Katze MG. 1991. The integrity of the stem structure of human immunodeficiency virus type 1 Tat-responsive sequence \*RNA\* is required for the interaction with the interferon-induced 68,000-Mr protein kinase. *J. Virol.* 65:632-40

211. Roy S, Parkin NT, Rosen...42

223. Serizawa H, Conaway JW, Conaway RC. 1994. An oligomeric form of the large subunit of transcription factor (TF) IIE activates phosphorylation of the \*RNA\* polymerase II carboxyl-terminal domain by TFIIH. *J. Biol. Chem.* 269:20750-56

224. Sheridan PL, Mayall TP, Verdin E, Jones KA. 1997. Histone acetyltransferases...

...cells. *Biochem. Biophys. Res. Commun.* 203:623-30

237. Tabernero C, Zolotukhin AS, Bear J, Schneider R, Karsenty G, et al. 1997. Identification of an \*RNA\* sequence within an intracisternal-A particle element able to replace Rev-mediated posttranscriptional regulation of human immunodeficiency virus type 1. *J. Virol.* 71:95-101

238. Tabernero C, Zolotukhin AS, Valentin A, Pavlakis GN, Felber BK. 1996. The posttranscriptional control element of the simian retrovirus type 1 forms an extensive \*RNA\* secondary structure necessary for its function. *J. Virol.* 70:5998-11

239. Tang H, Gaietta GM, Fischer WH, Ellisman MH, Wong-Staal F. 1997. A...

...of type D retrovirus. *Science* 276:1412-15

240. Tang H, McDonald D, Middlesworth T, Hope T, Wong-Staal F. 1999. The carboxyl terminus of \*RNA\* helicase A contains a bidirectional nuclear transport domain. *Mol. Cell. Biol.* In press

241. Trkola A, Dragic T, Arthos J, Binley JM, Olson BW, et... Jones KA. 1998. A novel CDK9-associated C-type cyclin interacts directly with HIV-1 Tat and mediates its high-affinity, loop-specific binding to \*TAR\* \*RNA\*. *Cell* 92:451-62

256. Weissenhorn W, Wharton SA, Calder LJ, Earl PL, Moss E, et al. 1996. The ectodomain of HIV-1 env subunit...

...Cell. *Biol.* 17:6898-905

265. Xu Y, Reddy TR, Fischer WH, Wong-Staal F. 1996. A novel hnRNP specifically interacts with HIV-1 RRE \*RNA\*. *J. Biomed. Sci.* 3:82-91

266. Yang X, Gold MO, Tang DN, Lewis DE, Aguilar-Cordova E, et al. 1997. TAK, an HIV Tat...

...X, Herrmann CH, Rice AP. 1996. The human immunodeficiency virus Tat proteins specifically associate with TAK in vivo and require the

carboxyl-terminal domain of \*RNA\* polymerase II for function. J. Virol. 70:4576-84

268. Yin L, Braaten D, Luban J. 1998. Human immunodeficiency virus type 1 replication is modulated...

**4/3,K/12 (Item 3 from file: 98)**

DIALOG(R)File 98:General Sci Abs/Full-Text

(c) 2003 The HW Wilson Co. All rts. reserv.

03546274 H.W. WILSON RECORD NUMBER: BGS197046274 (USE FORMAT 7 FOR FULLTEXT)

**Intracellular antibodies (intrabodies) for gene therapy of infectious diseases.**

Rondon, Isaac J

Marasco, Wayne A

Annual Review of Microbiology (Annu Rev Microbiol) v. 51 ('97) p. 257-83

SPECIAL FEATURES: bibl il ISSN: 0066-4227

LANGUAGE: English

COUNTRY OF PUBLICATION: United States

WORD COUNT: 11098

(USE FORMAT 7 FOR FULLTEXT)

TEXT:

... will then limit the spread of the virus in the patient (3). Different modalities of intracellular immunization have been developed, broadly divided into two categories: \*RNA\*-based and protein-based suppressors. The \*RNA\*-based suppressors include antisense \*RNA\*, ribozymes (130), and \*RNA\* decoys (141). Antisense \*RNA\* and ribozymes hybridize to target viral transcripts to inactivate them. \*RNA\* decoys interact and sequester regulatory proteins (e.g. Tat, Rev) that are essential for virus replication. The \*RNA\*-based suppressors are limited to the cytoplasmic compartment. The protein-based suppressors include transdominant mutant proteins (91), suicide molecules (14), and intracellular antibodies (96). Transdominant... AAV virus, and infection with several primary isolates from HIV-1 patients was effectively blocked in transduced T-lymphocytes (26).

**TAT**

Tat is a regulatory \*RNA\*-binding protein essential in the life cycle of HIV-1. HIV-1 encodes two forms of Tat, encoded by one (72 amino acid) or two (86 amino acid) exons. Tat protein activates transcription by association with a stable \*RNA\* hairpin transactivating region (\*TAR\*) that forms in the 5'-untranslated leader of nascent viral transcripts and promotes efficient elongation of transcription in vivo (82, 92, 97). This Tat-mediated...

...between Tat and Tat-binding cellular proteins, some of which may be transcription factors themselves (79, 111), as well as cellular proteins bound to the \*TAR\* region (54, 97) (Figure 5). In addition, Tat protein has other roles. Tat can easily be taken up by cells growing in tissue culture; it21). NF-kB binding to the enhancer elements in the HIV-1 LTR can cause \*TAR\*-independent activation of viral transcripts (11, 87). Thus, Tat is likely to have both direct and indirect effects in the pathogenesis of HIV-1.

The...

...stably inhibiting HIV-1 replication in transduced cells from HIV-1-infected patients than was sFv 105 in analogous studies.

**REV**

Rev is another regulatory \*RNA\*-binding protein essential in the life cycle of HIV-1. Rev is encoded by multiply spliced viral mRNA and appears to function efficiently only in a multimeric form (112). The Rev protein interacts with a short \*RNA\* sequence in the envelope gene Rev Responsive Element (RRE) to control the export of late mRNA from the nucleus; this

allows expression of virus structural the early stages of HIV-1 infection. RT specifically reverse transcribes the viral ribonucleic acid (\*RNA\*) to DNA, needed for integration into the host chromosome (Figure 7).

The anti-HIV-1 activity of anti-RT intrabodies has been studied by two

...neutralizing in assays of enzyme activity. The authors proposed that intracellular binding of anti-RT Fabs may sterically hinder movement of the enzyme along the \*RNA\* template or otherwise disrupt RT secondary structure.

Although most of the inhibitory effect on HIV-1 replication was observed in cell lines expressing the Fab...

...studies or virion infectivity studies.

#### NUCLEOCAPSID

The inner core of most retroviruses is formed by a shell of capsid protein molecules (p24) surrounding the dimeric \*RNA\* genome in close association with approximately 2000 molecules of nucleocapsid protein (NC), 20-50 molecules of reverse transcriptase and integrase (27), as well as tRNAs, 5S \*RNA\*, 7S \*RNA\*, and ribosomal RNAs (37). The NC protein contains two highly conserved zinc finger motifs (6). In vitro studies have shown that the NC is required for genomic \*RNA\* dimerization, for correct encapsidation, for annealing of the tRNA primer to its complementary binding site (PBS) on the genomic \*RNA\*, and for the minus-strand DNA transfer during reverse transcription (117, 121). These observations indicate that the NC plays an essential role in the afferent...sFv intrabodies.

Figure 5 Strategy for inhibition of Tat-mediated transactivation of viral gene expression by anti-Tat sFv intrabodies. Left side, Tat binding to \*TAR\* \*RNA\* and other cellular factors are required for transactivation. Right side, anti-Tat sFv binding to Tat may inhibit transactivation by inhibition of Tat nuclear import...

...of intron containing viral mRNAs. Left side, Rev binding to the cis-acting RRE sequence in unspliced and singly spliced viral mRNA leads to cytoplasmic \*RNA\* transport, viral structural protein synthesis, virus assembly, and production of infectious virions. Right side, anti-Rev sFv binding to any of several critical epitopes on...L, Buonaguro FM, Giraldo G, Ensoli B. 1994. The human immunodeficiency virus type 1 Tat protein transactivates tumor necrosis factor beta gene expression through a \*TAR\*-like structure. J. Virol. 68:2677-82

22. Burton DR, Barbas CF III, Persson MAA, Koenig S, Chanock RM, Lerner RA. 1991. A large array...

...fragments. Hum. Gene Ther. 6:1561-73

44. Duke GM, Hoffman MA, Palmenberg AC. 1992. Sequence and structural elements that contribute to efficient encephalomyocarditis virus \*RNA\* translation. J. Virol. 66:1602-9

45. Dul JL, Argon Y. 1990. A single amino acid substitution in the variable region of the light chain...272:872-77

52. Frankhauser C, Izaurralde E, Adachi Y, Wingfield P, Laemmler UK. 1991. Specific complex of human immunodeficiency virus type 1 Rev and \*nucleolar\* B23 proteins: dissociation by the Rev response element. Mol. Cell. Biol. 11:567-75

53. Fritz CC, Zapp ML, Green MR. 1995. A human nucleoporin...

...specifically interacts with HIV Rev. Nature 376:530-33

54. Gatignol A, Buckler-White A, Beckhout B, Jeang K-T. 1991. Characterization of a human \*TAR\* \*RNA\*-binding protein that activates the HIV-1 LTR. Science 251:1597-600

55. Glockshuber R, Malia M, Pfitzinger I, Pluckthun A. 1990. A comparison of...

...76. Kalland K-H, Szilvay AM, Langhoff E, Kaukenes G. 1994. Subcellular

distribution of human immunodeficiency virus type 1 Rev and colocalization of Rev with \*RNA\* splicing factors in a speckled pattern in the nucleoplasm. J. Virol. 68:1475-85

77. Kaloff CR, Haas IG. 1995. Coordination of immunoglobulin chain folding...Tiley LS, McCarn DF, Rusche JR, Hauber J, Cullen BR. 1990. HIV-1 structural gene expression requires binding of the Rev trans-activator to its \*RNA\* target sequence. Cell 60:675-83

94. Marasco WA. 1995. Intracellular antibodies (intrabodies) as research reagents and therapeutic molecules for gene therapy. Immunotechnology 1:1...

...93

97. Marciniak RA, Garcia-Blanco MA, Sharp PA. 1990. Identification and characterization of a HeLa nuclear protein that specifically binds to the trans-activation response (\*TAR\*) element of human immunodeficiency virus. Proc. Natl. Acad. Sci. USA 87:3624-38

98. Marks JD, Griffiths AD, Malmqvist M, Clackson TP, Bye JM, Winter ...virus infection. N. Engl. J. Med. 328(5):327-35

116. Park H, Davies MV, Langland JO, Chang H-W, Nam YS, et al. 1994. \*TAR\* \*RNA\*-binding protein is an inhibitor of the interferon-induced protein kinase PKR. Proc. Natl. Acad. Sci. USA 91:4713-17

117. Peliska JA, Balasubramanian S...

...G, Darlix JL. 1988. Small finger protein of avian and murine retroviruses has nucleic acid annealing activity and positions the replication primer tRNA onto genomic \*RNA\*. EMBO J. 7:1777-83

122. Pugsley AP. 1989. In Protein Targeting, San Diego: Academic

123. Richardson JH, Marasco WA. 1995. Intracellular antibodies: development and...

...viral life cycle. J. Virol. 70:3392-400

134. Sheng N, Erickson-Viitranen S. 1994. Cleavage of p15 in vitro by HIV-1 protease is \*RNA\* dependent. J. Virol. 68:6207-14

135. Shukla RR, Kimmel PL, Kumar A. 1994. Human immunodeficiency virus type 1 Rev-responsive element \*RNA\* binds to host cell-specific proteins. J. Virol. 68:2224-29

136. Siomi H, Shida H, Maki M, Hatanaka M. 1990. Effects of a highly basic region of human immunodeficiency virus Tat protein on \*nucleolar\* localization. J. Virol. 64:1803-7

137. Smith AE. 1995. Viral vectors in gene therapy. Annu. Rev. Microbiol. 49:807-38

138. Sodroski J, Goh...

...binding and particle assembly. J. Virol. 63:3232-42

141. Sullenger BA, Gallardo HF, Ungers GE, Gilboa E. 1991. Analysis of trans-acting response decoy \*RNA\*-mediated inhibition of human immunodeficiency virus type 1 transactivation. J. Virol. 65(12):6811-16

142. Taviadoraki P, Benvenuto E, Trinca S, De Martinis D...against human immunodeficiency virus type 1. Proc. Natl. Acad. Sci. USA 90:4156-60

150. Weiss R, Teich N, Varmus H, Coffin J, eds. 1985. \*RNA\* Tumor Virus. Cold Spring Harbor, NY: Cold Spring Harbor Lab.

151. Westendorp MO, Li-Weber M, Frank RW, Krammer PH. 1994. Human immunodeficiency virus type...

4/3,K/13 (Item 1 from file: 266)

DIALOG(R) File 266:FEDRIP

Comp & dist by NTIS, Intl Copyright All Rights Res. All rts. reserv.

00299035

IDENTIFYING NO.: 5R01AI50492-02 AGENCY CODE: CRISP

**Modeling \*RNA\*-based HIV gene therapeutics in SCID-hu mice**

PRINCIPAL INVESTIGATOR: AKKINA, RAMESH K

ADDRESS: AKKINA@LAMARCOLSTATE.EDU COLORADO STATE UNIVERSITY COLLEGE OF BETERINARY MED & BIOL FORT COLLINS, CO 80523

PERFORMING ORG.: COLORADO STATE UNIVERSITY, FORT COLLINS, COLORADO

SPONSORING ORG.: NATIONAL INSTITUTE OF ALLERGY AND INFECTIOUS DISEASES

DATES: 2001/01/02 TO 2012/31/06 FY : 2003



#### **Modeling \*RNA\*-based HIV gene therapeutics in SCID-hu mice**

...SUMMARY: vector transduced hematopoietic progenitor cells. Several new exciting developments occurred recently in the areas of stem cell biology, lentiviral gene transfer vectors, ribozyme targeting, and \*RNA\*-based therapeutics and, therefore, the stage is currently set to achieve success. In the present proposal, we would like to exploit these new technologies and...

... Project 2) are interactive and complimentary to the objectives of accompanying interactive R01 proposal (Project 1) by J. Rossi entitled "Combinatorial use of anti-HIV \*RNA\*-based therapeutics." The specific objectives of our proposal are: 1) Determine the effect of retrovirally transduced pol III promoter driven \*nucleolar\*, nuclear and cytoplasm targeted anti-HIV ribozymes \*TAR\* and RBE decoys, either individually or in combination, on the lineage specific differentiation of CD34+ cells into macrophages in vitro and into thymocytes in vivo in the SCID-hu thy/liv grafts and investigate the mechanism of action of \*RNA\*-based therapeutics in differentiated cells. 2) Determine the in vivo protective effects of different anti-HIV-1 RNAs, individually and in combination, in SCID-hu...

**4/3,K/14 (Item 2 from file: 266)**

DIALOG(R)File 266:FEDRIP

Comp & dist by NTIS, Intl Copyright All Rights Res. All rts. reserv.

00296693

IDENTIFYING NO.: 5R01AI42552-06 AGENCY CODE: CRISP

#### **Combinatorial use of anti-HIV \*RNA\*-based therapeutics**

PRINCIPAL INVESTIGATOR: ROSSI, JOHN J.

ADDRESS: JROSSI@COH.ORG BECKMAN RES INST OF CITY OF HOPE 1450 EAST DUARTE ROAD DUARTE, CA 91010-3011

PERFORMING ORG.: BECKMAN RESEARCH INST OF CITY OF HOPE, DUARTE, CALIFORNIA

SPONSORING ORG.: NATIONAL INSTITUTE OF ALLERGY AND INFECTIOUS DISEASES

DATES: 2001/01/98 TO 2012/31/06 FY : 2003

#### **Combinatorial use of anti-HIV \*RNA\*-based therapeutics**

...SUMMARY: transcriptase and protease for the treatment of HIV-1 infection. Patients on HAART have witnessed rapid decreases in blood and tissue levels of HIV-1 \*RNA\* and overall positive outcomes. Despite the overall success of HAART, reservoirs of HIV-1 continue to replicate in patients on HAART, necessitating prolonged and perhaps...

... overall goal of this program is to genetically modify human cells for resistance to HIV-1 infection and replication. To this end a series of \*RNA\* based anti-HIV-1 agents have been developed and tested. These inhibitory RNAs target multiple steps in the viral life cycle as well as the...

...led to the development of Pol III expressed, chimeric RNAs that localize the inhibitory RNAs to the cytoplasm, nucleus or nucleolus. We have shown that \*nucleolar\* localization of an anti-HIV-1 ribozyme, \*TAR\* or RBE decoy provides potent inhibition of viral replication. By combining the composition and intracellular localization of this collection of antiviral RNAs, it should be possible to achieve long-term, synergistic inhibition of HIV-1 replication. Importantly, the use of combinatorial inhibitory \*RNA\* therapy in conjunction with HAART may provide additional synergy, allowing reduced dosing of HAART reagents. The overall objectives of this proposal are to test the...

... inhibitory RNAs and several HAART reagents. This program is part of an interactive R01 program (Project 1) with Dr. Ramesh Akkina (Project 2) entitled "Modeling \*RNA\*-based HIV gene therapeutics in the SCID-hu mouse." The Specific Aims of this proposal are: 1) construction of retroviral and lentiviral vectors harboring multiple...

... RNAs and HAART agents in blocking infectious spread of HIV-1 in cell culture; 5) transduction of primary CD34+ cells with the single and combinatorial \*RNA\* expressing vectors to monitor differentiation in culture and differentiation in the SCID-hu mouse model. This aim is to be carried out collaboratively with Dr...

...DESCRIPTORS: combination chemotherapy; drug interaction; gene therapy; gene expression; human tissue; immunofluorescence technique; AIDS therapy; HIV infection; disease /disorder model; multidrug resistance; in situ hybridization; virus \*RNA\*; zidovudine; \*RNA\* directed DNA polymerase; nonhuman therapy evaluation; tissue /cell culture; virus replication; human immunodeficiency virus 1; northern blotting; ribozyme; transfection /expression vector; SCID mouse; CD34 molecule

4/3,K/15 (Item 3 from file: 266)

DIALOG(R)File 266:FEDRIP

Comp & dist by NTIS, Intl Copyright All Rights Res. All rts. reserv.

00295463

IDENTIFYING NO.: 2R37AI29329-14 AGENCY CODE: CRISP

**Enhancing the Intracellular Functioning of HIV Ribozymes**

PRINCIPAL INVESTIGATOR: ROSSI, JOHN J

ADDRESS: JROSSI@COH.ORG BECKMAN RES INST OF CITY OF HOPE 1450 EAST DUARTE ROAD DUARTE, CA 91010-3011

PERFORMING ORG.: BECKMAN RESEARCH INST OF CITY OF HOPE, DUARTE, CALIFORNIA

SPONSORING ORG.: NATIONAL INSTITUTE OF ALLERGY AND INFECTIOUS DISEASES

DATES: 2007/03/96 TO 2006/30/07 FY : 2002

...SUMMARY: in and demonstrations of efficacy both in vitro and in vivo have taken place. Ribozymes have the attributes of site-specific cleavage of the target \*RNA\*, the potential for turning over multiple substrates, and general lack of toxicity since they do not encode proteins In order to attain the goal of using anti-HIV ribozymes for treatment of HIV infection, we have been exploring several important areas of ribozyme function. These include identifying optimal \*RNA\* target sites for ribozyme interaction, co-localization of ribozyme and HIV RNAs and optimizing ribozyme expression from the backbones of lentiviral and retroviral vectors. The...

... analyses of engraftment potential. This latter work will be carried out collaboratively with Dr. Bruce Torbett at the Scripps Research Institute.

2) Functional analysis of \*nucleolar\* trafficking of HIV RNAs and regulatory proteins and in vivo selection for anti-HIV \*nucleolar\* localized ribozymes. A \*nucleolar\* localized anti-HIV-1 ribozyme as well as \*nucleolar\* localized \*TAR\* an RJ3E decoys have provided potent inhibition of HIV-1 infection in cultured cells. We will extend our analyses of these constructs to primary CD34+ cells and T-lymphocytes as part of specific aim 1. In aim 2 further analyses of the functional role of HIV-1 \*RNA\* trafficking through the nucleolus will be explored. Specifically, we will investigate the possibility that 2'0-methyl covalent backbone modifications in the R region of HN-1 \*RNA\* are guided by a cellular small \*nucleolar\* \*RNA\*. We will engineer a small \*nucleolar\* \*RNA\* to misdirect 2'0 methylations to specific sites of protein interaction in the HIV \*TAR\* and RJ3E elements. Finally, an in vivo SELEX scheme will be developoc to select for new targets, both in HIV-1 and cellular RNAs using a randomized binding arm library of \*nucleolar\* localized ribozymes. The results of the work proposed in this aim should extend our knowledge of the functional role of HIV \*RNA\* and protein trafficking through the nucleolus, and provide new ribozymes and other inhibitory RNAs for inhibition of HIV-1 infection. The overall objective of this work is to enhance our understanding of HIV \*RNA\* localization via the use of ribozymes, and to identify the best ribozyme-HIV target strategies for future use in human gene therapy.

4/3,K/16 (Item 1 from file: 357)

DIALOG(R)File 357:Derwent Biotech Res.

0295759 DBR Accession No.: 2002-17606

**A \*nucleolar\* \*TAR\* decoy inhibitor of HIV-1 replication - vector-mediated gene transfer and expression in host cell and polymerase chain reaction for use in HIV virus-1 gene therapy**

AUTHOR: MICHIEZI A; LI S; ZAIA JA; ROSSI JJ

CORPORATE AFFILIATE: City Hope Natl Med Ctr City Hope Natl Med Ctr

CORPORATE SOURCE: Rossi JJ, City Hope Natl Med Ctr, Beckman Res Inst, Div Mol Biol, 1450 E Duarte Rd, Duarte, CA 91010 USA

ISSN: 0027-8424 CODEN: 0027-8424; PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AME; (2002) 99, 22, 14047-14052

LANGUAGE: English

**A \*nucleolar\* \*TAR\* decoy inhibitor of HIV-1 replication - vector-mediated gene transfer and expression in host cell and polymerase chain reaction for use in HIV virus-1...**

...ABSTRACT: critical regulatory factor in HIV-1 gene expression. It mediates the transactivation of transcription from the HIV-1 LTR by binding to the transactivation response (\*TAR\*) element in a complex with cyclin T1. Because of its critical and early role in HIV gene expression, Tat and its interaction with the \*TAR\* element constitute important therapeutic targets for the treatment of HIV-1 infection. Based on the known \*nucleolar\* localization properties of Tat, we constructed a chimeric small \*nucleolar\* \*RNA\*-*\*TAR\** decoy that localizes to the nucleoli of human cells and colocalizes in the nucleolus with a Tat-enhanced GFP fusion protein. When the chimeric \*RNA\* was stably expressed in human T lymphoblastoid CEM cells it potently inhibited HIV-1 replication. These results demonstrate that the \*nucleolar\* trafficking of Tat is critical for HIV-1 replication and suggests a role for the nucleolus in HIV-1 viral replication. (6 pages)

DESCRIPTORS: \*nucleolar\* \*RNA\*-transactivation response decoy-inhibitor, TAT-enhanced green fluorescent protein prep., vector-mediated gene transfer, expression in host cell, polymerase chain reaction, appl. HIV virus-1...

**4/3,K/17 (Item 1 from file: 149)**

DIALOG(R) File 149:TGG Health&Wellness DB(SM)

(c) 2003 The Gale Group. All rts. reserv.

01421140 SUPPLIER NUMBER: 14003767 (USE FORMAT 7 OR 9 FOR FULL TEXT)

**Molecular targets of gene transfer therapy for HIV infection.**

Buchsacher, Gary L., Jr.

JAMA, The Journal of the American Medical Association, v269, n22, p2880(7) June 9, 1993

PUBLICATION FORMAT: Magazine/Journal ISSN: 0098-7484 LANGUAGE: English

RECORD TYPE: Fulltext; Abstract TARGET AUDIENCE: Professional

WORD COUNT: 7338 LINE COUNT: 00611

... target specific aspects of HIV replication, basic aspects of the HIV replication cycle will be reviewed briefly.

**HIV AND THE RETROVIRUS REPLICATION CYCLE**

Retroviruses are \*RNA\* viruses that replicate through a DNA intermediate. Virions contain two identical, single-stranded \*RNA\* molecules of positive polarity surrounded by the viral nucleocapsid protein. The remaining core of the virus is composed of the capsid and matrix proteins. In...

...expression from the HIV LTR. (6) Tat mediates this effect by interacting with a segment of the R region in the 5' LTR, termed the "*\*TAR\** element" (for trans-activating response), of transcripts initiated from the promoter. (7) The \*TAR\* element has been shown to form a stable stem loop structure. (8) Tat, along with unidentified cellular protein(s), (9,10) interacts with the \*RNA\* stem loop and apparently prevents premature termination of initiated transcripts. (11) Additional mechanisms of action,

such as increasing the translation efficiency of transcripts, also have...

...of full-length and singly spliced transcripts. (13,14) As with the Tat protein, Rev mediates its effect by interacting with a segment of transcribed \*RNA\*. The target of Rev is a segment of \*RNA\*, termed the rre (for Rev-responsive element), that is contained within the env gene. The rre has extensive secondary structure; interaction of Rev with the...

...in a pH-independent manner. (19,20) Following the release of the viral nucleic acid into the cytoplasm by an unknown mechanism, the single-stranded \*RNA\* molecules serve as a template for reverse transcriptase to generate a double-stranded DNA copy of the genome. (21,22)

The DNA copy of the...of viral structural proteins and replication enzymes. Full-length transcripts are incorporated into virion cores, probably through the interaction of capsid proteins with segments of \*RNA\* known to be important for encapsidation of the viral \*RNA\* into virions. (25'27) As virions bud from cells, they obtain their viral glycoprotein-containing envelopes. Released particles are then able to initiate a new...of this strategy, currently the titer of vector virus produced is far too low to be of practical use. Alternatively, isolation of the cis-acting \*TAR\* element or rre and their placement on a heterologous vector (eg, a murine retrovirus vector) can be used to limit gene expression to HIV-infected...

...with one or more steps in HIV replication but do not result in the death of the infected cells. These include, for example, viral-specific \*RNA\* ribozymes (39) or dominant negative mutants of viral gene products.

#### Elimination of Infected Cells

In an effort to stimulate a more effective immune response against...

...inhibition of protein synthesis. Initial experiments used the luciferase gene (luc) as a marker for the DT-A gene expressed from a construct containing the \*TAR\* element. (38) Although gene expression was induced in the presence of Tat, the results indicated that, for these genes, additional control of gene expression might...in the sequestering of cellular or viral components used in virus replication. An example of this is the use of high levels of expression of \*TAR\* sequences, which are known to interact with Tat and an unknown cellular factor. A chimeric tRNA-\*TAR\* construct, which was expressed in a T-cell line at high levels using a doublecopy murine retroviral vector, has been used to interfere with HIV replication. (47) Expression of high levels of \*TAR\* element sequences did not appear to affect cell viability. Further studies indicated that the overexpressed \*TAR\* elements were acting as decoys for a Tat-transactivation protein complex. (48) Another group used a Tat-inducible construct to express multimerized \*TAR\* elements. They found that induced transcription of these \*TAR\* sequences could decrease expression of a marker gene expressed from the HIV LTR. (49) Using a similar strategy, rre sequences were overexpressed in experiments using a T-cell line. (50) When challenged with HIV, these cells inhibited replication, though not as efficiently as cells expressing the \*TAR\* elements.

Antisense \*RNA\* molecules (sequences complementary to viral \*RNA\* sequences) also have been investigated as possible antiviral reagents. In this approach, it is hoped that the antisense molecules will hybridize to viral transcripts and thereby disrupt replication, translation, or encapsidation into virion particles of viral \*RNA\* molecules. In one study, antisense \*RNA\* corresponding to the HIV-1 primer binding site or to a region just 5' to the primer binding site was used to interfere with HIV replication in a CD4-positive lymphocyte-derived cell line. (51) Antisense \*RNA\* directed against tat and rev sequences also has been used to inhibit HIV replication. (52) After a T-lymphocyte line expressing antisense tat \*RNA\* molecules was infected with HIV, viral inhibition was observed for 2 weeks when a high multiplicity of infection was used. When a low multiplicity was used, however, inhibition continued for at least 5 weeks. Another group also found antisense \*RNA\* overlapping the tat gene initiation codon to be inhibitory, but again the effect was observed to be transient. (53)

Recently, an adeno-associated virus vector was used to introduce an

antisense sequence into both human hematopoietic and nonhematopoietic cell lines. (54) The antisense \*RNA\* used was an LTR sequence corresponding to the \*TAR\* element in the 5' LTR and part of the polyadenylation signal in the 3' LTR. Antisense \*RNA\* was produced constitutively and did not result in any apparent toxicity. When challenged with HIV, cells containing the antisense construct contained less viral mRNA, suggesting \*RNA\* hybrids were being rapidly degraded. Human immunodeficiency virus replication was inhibited up to 94% to 98% for 20 days as measured by a reverse transcriptase...

...99%. If these promising results can be extended to primary cells, this strategy may be effective for slowing HIV spread in vivo.

Ribozymes are small \*RNA\* molecules capable of catalyzing cleavage of specific \*RNA\* sequences. The use of ribozymes that target HIV sequences and that therefore might destroy HIV \*RNA\* before it can be replicated, translated, or packaged has been investigated. In one study, (55) a ribozyme specific for gag transcripts was expressed in CD4-positive HeLa cells. Constitutive expression of the ribozyme was found to decrease virus replication as assayed by viral antigen (p24) and \*RNA\* levels when these cells were challenged with HIV. Other attempts to use ribozymes specific for U5 sequences (56) or other 5' leader sequences (57, 58) also resulted in a varying detectable decrease in HIV replication. In addition, a ribozyme directed against HIV-1 integrase gene \*RNA\*, when expressed in *Escherichia coli*, has been shown to cleave the target sequence and to prevent integrase gene expression? Further work designed to identify more...

...be in considerable molar excess for the effect to be maximal. It has been suggested that the mutant Tat inhibited wild-type Tat by preventing \*nucleolar\* localization, possibly through the formation of inactive heterodimers.

A Rev routant that interferes with wild-type Rev function also has been identified. (64) The mechanism...initiation and stabilizes elongation. Cell. 1989;59:283-292.

8. Berkhout B, Silverman RH, Jeang K. Tat transactivates the human immunodeficiency virus through a nascent \*RNA\* target. Cell. 1989;59:273-282.

9. Gagnon A, Buclder-White A, Berkhout B, Jeang K. Characterization of a human \*TAR\* \*RNA\*-binding protein that activates the HIV-1 LTR. Science. 1991;251:1597-1600.

10. Marciniak RA, Garcia-Blanco MA, Sharp PA. Identification and characterization of a HeLa nuclear protein that specifically binds to the transactivation response (\*TAR\*) element of human immunodeficiency virus. Proc Natl Acad Sci U S A. 1990;87:3624-3628.

11. Cullen BR. The HIV-1 tat protein: an \*RNA\* sequence-specific processivity factor? Cell. 1990; 63:655-657.

12. Cullen BR. Trans-activation of human immunodeficiency virus occurs via a bimodal mechanism. Cell. 1986;46:973-982.

13. Emerman M, Vazeux R, Peden K. The rev gene product of the human immunodeficiency virus affects envelope-specific \*RNA\* localization. Cell. 1989; 57:1155-1165.

14. Felber BK, Hadzopoulou-Cladaras M, Cladaras C, Copeland T, Pavlakos GN. Rev protein of human immunodeficiency virus type...

...187-194.

24. Cullen BR, Greene WC. Regulatory pathways governing HIV-1 replication. Cell. 1989;58:423-426.

25. Aldovini A, Young R. Mutations of \*RNA\* and protein sequences involved in human immunodeficiency virus type 1 packaging result in production of noninfectious virus. J Virol. 1990;64:1920-1926.

26. Clavel F, Orenstein JM. A mutant of human immunodeficiency virus with reduced \*RNA\* packaging and abnormal particle morphology. J Virol. 1990;64:5230-5234.

27. Lever A, Gottlinger H, Haseltine W, Sodroski J. Identification of a sequence required for efficient packaging of human immunodeficiency virus type 1 \*RNA\* into virions. J Virol. 1989;63:4085-4087.

28. Freed EO, Myers DJ, Risser R. Mutational analysis of the cleavage sequence of the human immunodeficiency...HIV-regulated diphtheria toxin A

cbain gene. Hnm Gene Ther. 1992; 3:461-469.

47. Sullenget BA, Galiardo HF, Ungers GE, Gilboa E. Overexpression of \*TAR\* sequences renders cells resistant to human immunodeficiency virus replication. Cell. 1990;63:601-608.

48. Sullenger BA, Gallardo HF, Ungers GE, Gilboa E. Analysis of trans-acting response decoy \*RNA\*-mediated inhibition of human immunodeficiency virus type 1 transactivation. J Virol. 1991;65:6811-6816.

49. Lisziewicz J, Rappaport J, Dhar R. Tat-regulated production of multimerized \*TAR\* \*RNA\* inhibits HIV-1 gene expression. New Biologist. 1991;3: 82-89.

50. Lee TC, Sullinger BA, Gallardo HF, Ungers GE, Gilboa E. Overexpression of RRE...

...van Brunschot A, Asad S, van der Elst I, Read SE, Berstein A. Inhibition of human immunodeficiency virus type 1 multiplication by antisense and sense \*RNA\* expression. J Virol. 1991; 65:5524-5530.

52. Sczakiel G, Oppenlander M, Rittner K, Pawlira M. Tat- and Rev-directed antisense \*RNA\* expression inhibits and abolishes replication of human immunodeficiency virus type 1: a temporal analysis. J Virol. 1992;66:5576-5581.

53. Rhodes A, James W. Inhibition of human immunodeficiency virus replication in cell culture by endogenously synthesized antisense \*RNA\*. J Gen Virol. 1990;71:1965-1974.

54. Chatterjee S, Johnson PR, Wong KK Jr. Dualtarget inhibition of HIV-1 in vitro by means of...

...to human immunodeficiency virus type 1 (HIV-1) infection in human CD4+ lymphocyte-derived cell lines conferred by using retroviral vectors expressing an HIV-1 \*RNA\*-specific ribozyme. J Virol. 1991;65:5531-5534.

58. Ojwang JO, Hampel A, Looney DJ, Wong-Staal F, Rappaport J. Inhibition of human immunodeficiency virus...

...Kubota S, Furuta R, Maki M, Hatanaka M. Inhibition of human immunodeficiency virus type 1 Rev function by a Rev mutant which interferes with nuclear/\*nucleolar\* localization of Rev. J Virol. 1992;66:2510-2513.

67. Trono D, Feinberg MB, Baltimore D. HIV-1 Gag mutants can dominantly interfere with the...

**4/3,X/18 (Item 2 from file: 149)**

DIALOG(R) File 149:TGG Health&Wellness DB(SM)

(c) 2003 The Gale Group. All rts. reserv.

01313399 SUPPLIER NUMBER: 11517856 (USE FORMAT 7 OR 9 FOR FULL TEXT)

**The AIDS virus: what we know and what we can do about it.**

Wong-Staal, Flossie

The Western Journal of Medicine, v155, n5, p481(7)

Nov,

1991

PUBLICATION FORMAT: Magazine/Journal ISSN: 0093-0415 LANGUAGE: English

RECORD TYPE: Fulltext; Abstract TARGET AUDIENCE: Professional

WORD COUNT: 4527 LINE COUNT: 00463

...ABSTRACT: as a discussion of how different viral characteristics might be exploited for therapy. HIV is a retrovirus, which means that the viral genes consist of \*RNA\*. Unlike some \*RNA\* viruses, however, the retrovirus must make a DNA copy of the genes before it can replicate. (This 'backward' synthesis of DNA from \*RNA\* is the source of the 'retro' nomenclature.) This DNA copy of the viral genes is actually integrated directly into the chromosomes of the human host...

...replication or halting its spread. Halting the replication of the virus may focus on the viral enzyme known as reverse transcriptase, which transcribes the viral \*RNA\* into DNA. This approach benefits by the fact that no such enzyme exists in human cells under normal circumstances, and therefore a drug might inhibit...

... repeat (LTR). Infection begins with the binding of the viral

envelop to the cellular receptor, which then allows entry of the viral core containing the \*RNA\* genetic information. A viral enzyme, reverse transcriptase, converts the \*RNA\* to DNA provirus, which can then move to the nucleus, where it is integrated into the host chromosomal DNA, catalyzed by another viral enzyme, integrase...

...host chromosome, the infection is established and cannot be cured; only viral expression can be suppressed. The integrated provirus serves as a template for messenger \*RNA\* (mRNA) that can then be processed and translated into precursor proteins. The proteins then assemble and incorporate the genomic \*RNA\* to form virion particles. The maturation of these particles requires processing of the precursor proteins, using yet another viral enzyme, protease.

Every step in this...nucleus. The Rev protein, by interacting with a specific target sequence contained in the intron of the viral precursor mRNA, is able to disengage the \*RNA\* from the spliceosome and facilitate export into the cytoplasm (Figure 3). By doing so, Rev actually diversifies the distribution of multiply spliced \*RNA\* to unspliced or less spliced forms, so that Rev functions at the expense of its own production and the production of the other regulatory proteins...

...to this protein, all of them resulting in the activation of gene expression from the viral LTR. The target sequence that tat recognizes is called \*TAR\*, located immediately downstream of the site of transcription initiation. The tat gene has been known to increase the frequency of transcriptional initiation and to act...

...mechanisms, tat and rev also have several parallel features. They are both small nucleoproteins that are found predominantly in the nucleolus. Both of them recognize \*RNA\* rather than DNA targets, a feature that distinguishes them from most other viral transactivators. The target for rev is called the rev response element, or RRE, and the target for tat is \*TAR\*. In both cases, the \*RNA\* targets are highly structured molecules, and the recognition of them by the viral transactivators is largely structural. The rev gene product binds to a specific "hammerhead" (stem-loop II) region of the RRE, and tat binds to three unpaired bases in the so-called bulge of \*TAR\*. That binding of these viral transactivators to their target RNAs is not sufficient for function and that cellular factors are also involved are evident because...

...may not be functional. For example, the domain for binding to RRE has been mapped on the Rev protein. This region also doubles as the \*nucleolar\* targeting domain. In addition, there is another critical domain in which mutations do not affect the capability of the protein to bind to RRE, and ...

...that this region is an activator domain that mediates interaction of the Rev protein with cellular factor(s). Essential functional domains including the regions for \*RNA\* binding ...also what its role is in eukaryotic gene regulation.

Similarly, not one but several cellular factors have been shown to bind to either Tat or \*TAR\*. For example, a 68-kilodalton (kd) protein that recognizes the loop sequence of the \*TAR\* region has been described. A 45-kd protein with the same \*TAR\*-binding specificity is prevalent in T cells. A similar-sized protein binds specifically to the stem between the loop and the bulge (Figure 4). Again...

...by expression of the core protein p24 or viral reverse transcriptase. The sense-strand of the same sequence had no effect. When the expression of \*RNA\* is examined in the antisense-treated cells, neither tat nor rev is completely inhibited--that is, some \*RNA\*, including singly spliced viral \*RNA\*, is expressed. The unspliced genomic \*RNA\*, however (the gag-pol message), appears to be highly sensitive to rev depletion and is completely repressed. As a result, viral production is also repressed...  
...gene an attractive target for virus inhibition.

As discussed earlier, both tat and rev transactivations involve a number of players: the viral protein, a viral \*RNA\* element, and cellular factors, each interacting with the others. It is possible to devise means

of interrupting those interactions that are crucial to their functions. For example, some mutant proteins are negative dominant. These proteins can retain binding activity to their \*RNA\* targets but are not able to interact with the proper cellular factors and therefore would not function. They can thus act as competitive inhibitors of the wild-type protein. Alternatively, oligonucleotides that can hybridize to the stem-loop sequences of the \*RNA\* target can presumably disrupt the structure or the folding of these molecules, thereby rendering them incapable of interacting with both viral and cellular factors. Synthetic...Klotman, S. Daefler, F. Wong-Staal, unpublished data, 1991).

There are many variations on this theme. One example is the expression of multiple copies of \*TAR\* \*RNA\* as a decoy to compete with normal \*TAR\* in the virus genome for expression. A potential problem of expressing a \*TAR\* or RRE decoy, however, is that it may also sequester cellular factors that may be important for cell function. To overcome this problem, a vector has been devised in which the expression of \*TAR\* is conditional on tat acting on its promoter, which is the HIV LTR. Therefore, these decoy molecules would only be expressed in infected cells, where the first sign of the expression of tat can then unleash a whole army of these \*TAR\* decoys to limit the extent of viral replication (J. Lisiewicz, MD, written communication, April 1991).

The foregoing discussion only relates to laboratory successes on inhibiting...

**4/3,K/19 (Item 3 from file: 149)**

DIALOG(R)File 149:TGG Health&Wellness DB(SM)

(c) 2003 The Gale Group. All rts. reserv.

01311360 SUPPLIER NUMBER: 10873149 (USE FORMAT 7 OR 9 FOR FULL TEXT)

**Arginine-mediated \*RNA\* recognition: the arginine fork.**

Calnan, Barbara J.; Tidor, Bruce; Biancalana, Sara; Hudson, Derek; Frankel, Alan D.

Science, v252, n5009, p1167(5)

May 24,

1991

PUBLICATION FORMAT: Magazine/Journal ISSN: 0036-8075 LANGUAGE: English

RECORD TYPE: Fulltext TARGET AUDIENCE: Academic

WORD COUNT: 3418 LINE COUNT: 00325

**Arginine-mediated \*RNA\* recognition: the arginine fork.**

TEXT:

...ARE important for many regulatory processes, but little is known about the details of sequence-specific recognition. From what is known, it appears that both \*RNA\* structure and nucleotide sequence function in recognition. The crystal structure of the glutamyl tRNA synthetase-tRNA complex [1] has shown that specific contacts are made between amino acid side chains and bases in non-base paired regions of the \*RNA\*, while studies of the R17 coat protein [2] have suggested that the overall three-dimensional \*RNA\* conformation contributes substantially to recognition. Recently, an arginine-rich \*RNA\*-binding motif has been identified in several \*RNA\*-binding proteins [3], including the human immunodeficiency virus (HIV) Tat protein. Peptides that contain this region of Tat bind specifically to an \*RNA\* stem-loop structure named \*TAR\* [4, 5], which is located in the HIV long terminal repeat, and \*RNA\* binding is essential for Tat-dependent transcriptional activation [5]. The overall charge density of the Tat peptides is important for binding, however, the amino acid sequence requirements are flexible; the sequence can be scrambled and still bind specifically to \*TAR\* [5].

The basic \*RNA\*-binding region of Tat, RKRRQRRR (residues 49 to 57), is nine amino acids long and contains a glutamine at position 54 that is not essential for binding or activity [5]. Because it is known that a high positive charge density is important for \*RNA\* binding, we synthesized [6] two peptides, [R.sup.9], which contains a stretch of nine adjacent arginines (with a tyrosine at the [NH.sub.2]...



...the COOH-terminus), and [K.sub.9], which contains a stretch of nine lysines (and a surrounding tyrosine and alanine), and measured their binding to \*TAR\* \*RNA\* [7]. The [R.sub.9] peptide bound to \*TAR\* \*RNA\* with the same affinity as the wild-type Tat peptide and with ten-fold higher affinity than [K.sub.9] (Fig. 1). The specificity of [R.sub.9] binding to \*TAR\* was identical to the wild-type peptide, whereas [K.sub.9] binding was nonspecific [7]. Because \*RNA\* binding of Tat peptides correlates with Tat's function as a transcriptional activator [5], we asked whether [R.sub.9] or [K.sub.9] could...

...and was 100-fold more active than the [K.sub.9]-containing protein (Fig. 2). These results confirm that transactivation of transcription correlates with peptide \*RNA\* binding and suggest that arginine residues are important for specific \*RNA\* recognition.

The low level of transactivation achieved by the [K.sub.9]-containing protein allowed us to systematically replace lysines with arginines in order to...

...52 to 53 restored transactivation to wild-type levels (Fig. 3). Electrophoretic mobility shift experiments with YKKKRKKKKKA (R52) or with YKKKRKKKKKA (R53) peptides showed that \*RNA\*-binding affinity and specificity were also restored to wild-type levels by the addition of one arginine at either position 52 or 53 [10]. To...

...basic region (Fig. 4). Thus, a single arginine surrounded by three to four basic amino acids on each side is sufficient for specific recognition of \*TAR\* \*RNA\*.

These and other results [4, 5, 11] clearly suggested that \*RNA\* structure is important in Tat-\*TAR\* recognition. It seemed plausible that the \*RNA\* backbone might be adopting a highly defined conformation and that a specific configuration of phosphates was being recognized by arginine. To identify phosphates involved in...

...contacts two adjacent phosphates (even in the wild-type peptide, which contains multiple arginines), although we cannot rule out the possibility that ethylation alters the \*RNA\* structure and indirectly interferes with binding. Ethylation of some phosphates, particularly the phosphate between G26 and A27, seems to enhance binding (Fig. 5), possibly by stabilizing the \*RNA\* structure. The minimal interference observed with lysines or with nonspecific arginines suggests that these residues may make weaker contacts with the \*RNA\* or that alternative phosphates may be contacted when one is modified.

How does a single arginine recognize \*TAR\* \*RNA\*? Arginine contains two terminal amino ([NH.sub.2]) groups at the [eta] position and a secondary amine (NH) at the [epsilon] position, each of which...

...and geometries are shown in Fig. 6A. Clearly, an arginine side chain can form many possible hydrogen bonds with appropriately positioned acceptor groups on the \*RNA\*. These acceptors can include phosphate oxygens, the ribose 2' OH, and groups on the bases (for example, O-6 and N-7 on guanine or...

...the planar geometry of the arginine amino groups. Our ethylation interference data indicate that a single arginine in Tat contacts two adjacent phosphates at the \*TAR\* bulge, suggesting that the phosphate backbone adopts a defined conformation that can be bridged by arginine in a fork-like arrangement. To determine a plausible...phosphorus atoms). We define the arginine fork as an interaction between a single arginine and a pair of adjacent phosphates, which mediates specific recognition of \*RNA\* structure. Other arginine-phosphate arrangements are possible (for example, see legend to Fig. 6B), and arginine forks with additional H-bonds are possible (for example, with a specific base or a 2' OH).

To determine whether such phosphate arrangements are found in \*RNA\* structures, the modeled phosphate coordinates from Fig. 6B were superimposed on all phosphate pairs in tRNA crystal structures [14]. The results indicate that double-stranded A-form \*RNA\* cannot readily accommodate this arrangement; the P-P distance in the model (7.1

[angstroms]) is longer than the P-P distance in A-form \*RNA\* (5.6 [angstroms]), and the phosphate oxygens in A-form \*RNA\* are not properly oriented to form H-bonds between a single arginine and a pair of adjacent phosphates. Reasonable H-bonding arrangements are much more likely to be found at discontinuous regions of \*RNA\*, for example, at junctions between double-stranded A-form \*RNA\* and a bulge or loop. The two critical phosphates in \*TAR\* are located precisely at the junction of the double-stranded stem and the 3-nucleotide bulge.

The cocrystal structure of glutamyl tRNA synthetase-tRNA shows...

...additional H-bond with a ribose 2' OH. It is plausible that the arginine in Tat also interacts with a 2' OH, thus discriminating between \*RNA\* and DNA [4]. Although we cannot rule out base-specific contacts, for example, with an essential uridine in the bulge [4, 5, 11], it seems...

...contacts between one arginine and a highly oriented pair of phosphates can account for the modest 10- to 20-fold specificity of Tat binding to \*TAR\* [4, 5, 11]. The interactions that stabilize the overall folding of \*TAR\* and the orientation of these particular phosphates remain to be determined. In addition to the specific arginine contact, the charge density of the basic region...

...Tat is important for binding [5] and may provide a nonspecific electrostatic scaffold to help orient the arginine. Tat is perhaps the simplest example of \*RNA\* recognition in that a single amino acid interacts with a single feature of the \*RNA\*; other proteins may achieve higher specificity through multiple arginine-\*RNA\* or other interactions. In the case of Tat, although \*RNA\* binding is essential for transactivation, the modest specificity for \*TAR\* is insufficient to account for the high specificity of Tat function. Other interactions of Tat, perhaps with cellular proteins, are likely to be required.

The recognition of \*TAR\* by Tat highlights fundamental differences between \*RNA\* recognition and DNA recognition. It is clear from the structures of protein-DNA complexes that sequence-specific discrimination derives primarily from direct base-specific contacts...

...commonly made in the DNA major groove [5]. In most cases, DNA tertiary structure does not seem to be of major importance in recognition. That \*RNA\* recognition often seems to rely on \*RNA\* tertiary structure is emphasized by the finding that the Tat-\*TAR\* interaction uses only a single arginine side chain in the midst of an apparently unstructured segment of basic amino acids [5] to recognize a specific backbone conformation of \*TAR\*. The unstructured nature of the unbound polypeptide is supported by the fact that the sequence of the Tat basic region can be simplified to a single arginine embedded in a set of eight lysines. The peptide conformation when bound to \*TAR\* remains to be determined. Specific recognition of \*TAR\* appears to occur by indirect readout of the base sequence and direct contact with the phosphate backbone, and may not involve any base-specific contacts. Differences between DNA and \*RNA\* recognition are also apparent in studies of TFIIIA, a zinc finger-containing protein that binds to the same site on both DNA and \*RNA\*. DNA recognition seems to occur through base-specific contacts in the major groove; \*RNA\* recognition appears to be primarily backbone structure-specific [16].

Other \*RNA\* hairpins and bulges form stable, ordered tertiary structures [17], further emphasizing that \*RNA\* structure can provide information for protein-\*RNA\* complex formation. The structure of an \*RNA\* pseudoknot reveals that phosphates can be arrayed in unusual, electrostatically unfavorable geometries through tertiary \*RNA\* interactions [18]; binding of basic amino acids to these phosphates might help to stabilize a more favorable \*RNA\* conformation that, in turn, would provide favorable energy for the protein-\*RNA\* interaction. This could explain why interaction of HIV Tat or Rev with \*RNA\* causes a change in \*RNA\* conformation upon binding [5, 19].

Other \*RNA\*-protein interactions will likely follow some of the principles outlined here. Arginine-rich motifs similar to the basic region of Tat are found in several \*RNA\*-binding proteins, including bacterial

antiterminators, ribosomal proteins, and HIV Rev [3]. Other \*RNA\*-binding proteins may bring basic amino acids together through protein tertiary structure rather than primary sequence and may position specific arginines to interact with defined \*RNA\* structures. For example, the U1 A protein, which contains a ribonuclear protein (RNP) \*RNA\*-binding motif, has a highly defined structure with a cluster of basic amino acids at one end and at least one arginine that is essential...

...recognition [20]. It is not yet known if this arginine makes a base-specific or structure-specific contact. For TFIIIA, the strongest interactions with 5S \*RNA\* are localized to junctions between stems and loops [21], similar to the stem-bulge junction in \*TAR\*; perhaps arginines participate in some of these interactions. Arginine has also been shown to bind to the guanosine binding site of a group I intron [22]; however, this interaction involves H-bonding to a guanine base in the \*RNA\* and is distinct from the arginine fork proposed here. Many \*RNA\*-binding proteins, including heterogeneous nuclear \*RNA\*-binding proteins and \*nucleolar\* proteins, contain clusters of methylated arginines, most commonly [N.sup.G], [N.sup.G]-dimethylarginine [23]. Because methylation would block H-bonding but would not alter the charge of the side chain, arginine methylation could provide a mechanism to regulate \*RNA\* binding between specific and nonspecific modes. While it is clear the \*RNA\* recognition will involve more than just arginine forks, it seems reasonable to suggest that arginine-mediated recognition of \*RNA\* structure may be an important part of many \*RNA\*-protein complexes.

#### REFERENCES AND NOTES

- [1] M. A. Rould, J. J. Perone, D. Soll, T. A. Steitz, *Science*, 246, 1135 (1989).
- [2] P. J. Romaniuk...

...spectrometry (University of California, Berkeley), and purity and concentrations were confirmed by native polyacrylamide (20%) gel electrophoresis in sodium acetate, pH 4.5 (30mM).

[7] \*TAR\* RNAs were transcribed by T7 \*RNA\* polymerase in vitro with synthetic oligonucleotide templates [5] [J. F. Milligan and O. C. Uhlenbeck, *Methods Enzymol.* 180, 51 (1989)]. All RNAs contained GG at their 5' end, which increased the efficiency of transcription, and CC at the 3' end to base pair with GG. Wild-type \*TAR\* \*RNA\* contained sequences +18 to +44 of the HIV long terminal repeat \*TAR\* site. Mutant TARs were also synthesized and specificity of binding of each peptide was determined [5]. All RNAs were purified on 10% polyacrylamide/8 M...

...in ammonium acetate (0.5 M), magnesium acetate (10 mM), EDTA (1 mM), and SDS (0.1%), extracted twice with phenol, and ethanol-precipitated. Purified \*RNA\* was resuspended in sterile deionized water. The concentrations of labeled RNAs were determined from the specific activity [sub.32.P]CTP incorporated into the transcripts. Unlabeled \*RNA\* was quantitated by spectrophotometry. \*RNA\* electrophoretic mobility shift assays were performed as described [5]. Briefly, peptide and \*RNA\* were incubated together for 10 min on ice in binding reactions (10 [microliter]) that contained tris-HCl pH 7.5 (10 mM), NaCl (70 mM), EDTA (0.2 mM), and glycerol (5%). Peptide-\*RNA\* complexes were resolved on polyacrylamide (10%) gels in tris-borate-EDTA [0.5 X TBE; in *Current Protocols in Molecular Biology*, F. M. Ausubel et...

...U. Delling, C.-H. Chen, C. A. Rosen, N. Sonenberg, *Genes Dev.* 4, 1365 (1990); C. Dingwall et al., *EMBO J.* 9, 4145 (1990).

[12] \*TAR\* \*RNA\* (31 nucleotides) was labeled with [sub.32.P] at the 5' end with T4 polynucleotide kinase. \*RNA\* phosphates were ethylnitrosourea under denaturing conditions as described [V. V. Vlassov, R. Giege, J.-P. Ebel, *Eur. J. Biochem.* 119, 51 (1981)], except...

...using a saturated solution of ethylnitrosourea in ethanol) and tRNA (2 [microgram], and were incubated at 80 [degrees] C for 5 min. After modification, the \*RNA\* was ethanol-precipitated, washed with 100% ethanol and lyophilized. Peptides were bound to the modified \*RNA\* (500,000 cpm) at concentrations that gave < 50% binding (specific) by gel shift [7] or at

higher concentrations that gave nonspecific binding. Free and bound RNAs were visualized by autoradiography. The bands were excised and the \*RNA\* was eluted from the gel [7], ethanol precipitated with yeast tRNA (20 [microgram]), and lyophilized. To cleave the phosphotriester bonds (at the modified phosphates), samples...

...i+2), and those distant in primary sequence but near each other in the tertiary structure. In no case did phosphate pairs in double-stranded \*RNA\* match the template. The (i, i+1) pattern frequently appeared surrounding the first or last unpaired base in a bulge or loop, and the (i...

...DESCRIPTORS: \*RNA\*--

**4/3,K/20 (Item 1 from file: 159)**  
DIALOG(R)File 159: Cancerlit  
(c) format only 2002 Dialog Corporation. All rts. reserv.

01915587 PMID: 93686277

**HIV-1 transactivator.**

No affiliation given

Harvard AIDS Inst Ser 1991, 1 p43-310,

Document Type: MONOGRAPH

Languages: ENGLISH

Main Citation Owner: NOTNLM

Record type: Completed

...viral regulatory proteins tat and rev, are expressed, whereas, late in infection production switches to full-length, unspliced transcripts that act both as the virion \*RNA\* and the mRNA for the gag-pol polyprotein. HIV-1 transactivator is reviewed in the following chapters: \*RNA\* interactions of the tat and rev proteins of HIV-1; the role of the \*TAR\* region and tat protein in HIV-directed gene expression; \*RNA\*-protein interactions required for regulation of HIV gene expression; cellular factors involved in regulating HIV gene expression; biochemical characterization of REV-RRE interaction; analysis of...

... the rex protein of HTLV-1; tat-mediated trans-activation as a presplicing event requiring a functional HIV-1 TATAA element; analysis of the TAT/\*TAR\* interaction in Xenopus oocytes; transcriptional regulation of HIV; functional topography of lentivirus tat proteins defined by domain switching; \*nucleolar\* targeting singles of HIV; characterization of the specific interaction between TAT and \*TAR\* \*RNA\* ; and the exogenous trans-activation activity of HIV-1 tat.

Chemical Name: Gene Products, rev; Gene Products, rex; Gene Products, tat ; \*RNA\*, Viral; Trans-Activators; Viral Proteins

**4/3,K/21 (Item 1 from file: 444)**  
DIALOG(R)File 444: New England Journal of Med.  
(c) 2003 Mass. Med. Soc. All rts. reserv.

00110787

Copyright 1992 by the Massachusetts Medical Society

**Medical Progress: Nonmelanoma Cancers Of The Skin (Review Article)**

Preston, Diana; Stern, Robert.

The New England Journal of Medicine

Dec 3, 1992; 327 (23), pp 1649-1662

LINE COUNT: 00688

WORD COUNT: 09497

**TEXT**

...58-74). In 1775, Sir Percival Pott reported an increased risk of squamous-cell carcinoma of the genitals in English chimney sweeps exposed to coal-\*tar\* products, the first recognized example of a chemical carcinogen (Ref. 75). Numerous chemicals are associated with an increased

risk of nonmelanoma skin cancer, primarily squamous...deeper tumors (Ref. 153-155). A fluorinated pyrimidine, fluorouracil inhibits the methylation of deoxyuridylic acid to thymidylic acid, interfering with the synthesis of DNA and \*RNA\* and cell growth (Ref. 156). Available as creams and solutions, fluorouracil is applied twice

#### CITED REFERENCES

- ...tumor infiltration in basal cell carcinoma. J Dermatol Surg Oncol 1991;17:574-8.
102. Kanitakis J, Hoyo E, Hermier C, Chouvet B, Thivolet J. \*Nucleolar\* organizer region enumeration in keratoacanthoma and squamous cell carcinomas of the skin. Cancer 1992;69:2937-41.
103. Sanders GH, Miller TA. Are keratoacanthomas really...

4/3,K/22 (Item 2 from file: 444)

DIALOG(R)File 444:New England Journal of Med.

(c) 2003 Mass. Med. Soc. All rts. reserv.

00108320

Copyright 1991 by the Massachusetts Medical Society

#### **Mechanisms of Disease: The Molecular Biology Of Human Immunodeficiency Virus Type 1 Infection (Review Article)**

Greene, Warner C.

The New England Journal of Medicine

Jan 31, 1991; 324 (5),pp 308-317

LINE COUNT: 00566

WORD COUNT: 07820

#### TEXT

...whereas the pol gene gives rise to the viral reverse transcriptase and other enzymatic activities (Fig. 1). HIV-1, however, contains in its 9-kilobase \*RNA\* genome not only these three essential genes but also at least six additional genes (vif, vpu, vpr, tat, rev, and nef) (Fig. 1). It is...

...3' long terminal repeats (LTRs) containing regulatory sequences recognized by various host transcription factors are also depicted, and the positions of the Tat and Rev \*RNA\* response elements (\*TAR\* transactivation response] element and Rev response element) are indicated \*. \*\*FIGURE OMITTED...inner surface of the lipid bilayer and probably stabilizes the exterior and interior components of the virion. The p7 protein binds directly to the genomic \*RNA\* through a zinc-finger structural motif and together with p9 forms the nucleoid core. Importantly, this retroviral core also contains two copies of the single-stranded HIV-1 genomic \*RNA\* that is associated with the various preformed viral enzymes, including the reverse transcriptase, integrase, and protease (Fig. 2).

\*Figure 2.-Schematic Diagram of the HIV...

...and gp41(sup env)) and nucleocapsid (p24(sup gag), p17(sup gag), p9(sup gag), and p7(sup gag)) is identified. In addition, the diploid \*RNA\* genome is shown associated with reverse transcriptase, an \*RNA\*-dependent DNA polymerase \*. \*\*FIGURE OMITTED...

...the entry of HIV-1 into the cell (lck denotes a lymphoid-specific tyrosine kinase that binds to CD4). After uncoating, reverse transcription of viral \*RNA\* begins, resulting in the production of the double-stranded DNA form of the virus in the presence of the appropriate host factors. In turn, the...for the replicative phase of its life cycle (Ref. 6). Viral replication begins with the generation of a first-strand DNA copy of the viral \*RNA\* mediated by the HIV-1-encoded reverse transcriptase (Fig. 3). Second-strand DNA synthesis is also controlled by the reverse transcriptase but proceeds only after the action of a second pol-gene product, ribonuclease H, which partially degrades the original \*RNA\* template. When complete, the reverse-transcription reaction yields a double-stranded DNA replica of the original \*RNA\* genome containing tandem long terminal repeats at each end of the DNA in lieu of the short terminal repeats